

Geometric Bäcklund–Darboux transformations for the KP hierarchy

G.F. Helminck

J.W. van de Leur*

Faculty of Applied Mathematics,
University of Twente,
P.O.Box 217, 7500 AE Enschede,
The Netherlands
fax: 31-53 489 4824
e-mail: helminck@math.utwente.nl
vdleur@math.utwente.nl

February 4, 2008

1991 MSC : 22E65, 22E70, 35Q53, 35Q58, 58B25

Keywords : KP hierarchy, Bäcklund-Darboux transformation,
Grassmann manifold

Abstract

In this paper it is shown that, if you have two planes in the Segal-Wilson Grassmannian that have an intersection of finite codimension, then the corresponding solutions of the KP hierarchy are linked by Bäcklund-Darboux (shortly BD-)transformations. The pseudodifferential operator that performs this transformation is shown to be built up in a geometric way from so-called elementary BD-transformations and is given here in a closed form. The geometric description of elementary BD-transformations requires that one has a geometric interpretation of the dual wavefunctions involved. This is done here with the help of a suitable algebraic characterization of the wavefunction. The BD-transformations also induce transformations of the tau-function associated to a plane in the Grassmannian. They are determined explicitly for the elementary and so-called trivial BD-transformations. For the Gelfand-Dickey hierarchies we derive a geometric characterization of the BD-transformations that preserves these subsystems of the KP hierarchy. This generalizes the classical Darboux-transformations. We also determine an explicit expression for the squared eigenfunction potentials.

Next a connection is laid between the KP hierarchy and the 1-Toda lattice hierarchy. It is shown that infinite flags in the Grassmannian yield solutions of the latter hierarchy. These flags can be constructed by means of BD-transformations, starting from some plane. Other applications of these BD-transformations are a geometric way to characterize Wronskian solutions of the m -vector k -constrained KP hierarchy and the construction of a vast collection of orthogonal polynomials that are parametrized among other things by the commuting flows of the KP hierarchy. These systems of polynomials play a role in matrix models.

*JvdL is financially supported by the Netherlands Organization for Scientific Research (NWO).

1 Introduction to the KP Hierarchy

The KP hierarchy consists of a tower of nonlinear differential equations in infinitely many variables $\{t_n | n \geq 1\}$. It is named after the simplest nontrivial equation in this tower, the Kadomtsev–Petviashvili equation:

$$\frac{3}{4} \frac{\partial^2 u}{\partial t_2^2} = \frac{\partial}{\partial t_1} \left(\frac{\partial u}{\partial t_3} - 3u \frac{\partial u}{\partial t_1} - \frac{1}{4} \frac{\partial^3 u}{\partial t_1^3} \right). \quad (1.1)$$

The solutions of these equations belong to a ring of functions R that is stable under the operators $\partial_n = \frac{\partial}{\partial t_n}$. The form in which one presents these equations, is the so-called Lax form. This is an equality between operators in the privileged derivation $\partial = \partial_1$ of a specific nature. For notational convenience we sometimes write x instead of t_1 and $\frac{\partial}{\partial x}$ instead of ∂ . The simplest way to present these equations, is to extend the ring $R[\partial] = \{\sum_{i=0}^n a_i \partial^i | a_i \in R\}$ by adding suitable integral operators to the ring. Thus it becomes possible to take the inverse and roots of certain differential operators. Consider the ring $R[\partial, \partial^{-1}]$ of pseudodifferential operators with coefficients in R . It consists of all expressions

$$\sum_{i=-\infty}^N a_i \partial^i, \quad a_i \in R \quad \text{for all } i,$$

that are added in an obvious way and multiplied according to

$$\partial^j \circ a \partial^i = \sum_{k=0}^{\infty} \binom{j}{k} \partial^k(a) \partial^{i+j-k}.$$

Each operator $P = \sum p_j \partial^j$ decomposes as $P = P_+ + P_-$ with $P_+ = \sum_{j \geq 0} p_j \partial^j$ its differential operator part and $P_- = \sum_{j < 0} p_j \partial^j$ its integral operator part. We denote by $\text{Res}_\partial P = p_{-1}$ the *residue* of P .

An operator $L \in R[\partial, \partial^{-1}]$ of the form

$$L = \partial + \sum_{j < 0} \ell_j \partial^j, \quad \ell_j \in R \quad \text{for all } j < 0. \quad (1.2)$$

is called a *Lax operator*. We say that a Lax operator satisfies the KP hierarchy if and only if it satisfies the system of equations

$$\partial_n(L) = \sum_{j < 0} \partial_n(\ell_j) \partial^j = [(L^n)_+, L], \quad n \geq 1, \quad (1.3)$$

called the *Lax equations* for L . These Lax equations are equivalent to the following infinite set of equations, which are called the *Zakharov–Shabat equations*:

$$\partial_n(L^m)_+ - \partial_m(L^n)_+ = [(L^n)_+, (L^m)_+], \quad m, n \geq 1. \quad (1.4)$$

Hence both (1.3) and (1.4) can be seen as the compatibility equation of the linear system

$$L\psi = z\psi \quad \text{and} \quad \partial_n(\psi) = (L^n)_+(\psi) \quad (1.5)$$

Notice that the equation (1.3) has at least the trivial solution $L = \partial$ and for this solution (1.5) becomes

$$\partial\psi = z\psi \quad \text{and} \quad \partial_n\psi = z^n\psi \quad \text{for all } n \geq 1.$$

Hence, the function $\gamma(t) = \exp(\sum_{i \geq 1} t_i z^i)$ is a solution. The space M of so-called *oscillating functions* is a space for which we can make sense of (1.5) and that can be seen as a collection of perturbations of this solution. It is defined as

$$M = \{(\sum_{j \leq N} a_j z^j) e^{\sum t_i z^i} \mid a_i \in R, \text{ for all } i\}.$$

Expressing $e^{\sum t_i z^i}$ in terms of the elementary Schur functions, i.e.,

$$e^{\sum t_i z^i} = \sum_{k=0}^{\infty} p_k(t) z^k$$

one notices that the product

$$(\sum_{j \leq N} a_j z^j) e^{\sum t_i z^i} = \sum_{\ell \in \mathbb{Z}} (\sum_{k=0}^{\infty} a_{\ell-k} p_k) z^{\ell}$$

is still formal. To make sense of this expression as an infinite series in z and z^{-1} , the coefficients $\sum_{k=0}^{\infty} a_{\ell-k} p_k$ for all $\ell \in \mathbb{Z}$ have to be well defined functions of $t = (t_n)$. In that light it is natural to have a context such that for each element t

$$\sum_{j \leq N} |a_j(t)|^2 < \infty \quad \text{and} \quad \sum_{k=0}^{\infty} |p_k(t)|^2 < \infty.$$

Such a context has been given in [22] and will be considered in section 2.

The space M becomes a $R[\partial, \partial^{-1}]$ -module by the natural extension of the actions

$$\begin{aligned} b\{(\sum_j a_j z^j) e^{\sum t_i z^i}\} &= (\sum_j b a_j z^j) e^{\sum t_i z^i} \\ \partial\{(\sum_j a_j z^j) e^{\sum t_i z^i}\} &= (\sum_j \partial(a_j) z^j + \sum_j a_j z^{j+1}) e^{\sum t_i z^i}. \end{aligned}$$

It is even a free $R[\partial, \partial^{-1}]$ -module, since we have

$$(\sum p_j \partial^j) e^{\sum t_i z^i} = (\sum p_j z^j) e^{\sum t_i z^i}.$$

An element ψ in M is called an *oscillating function of type z^ℓ* , if it has the form

$$\psi(z) = \psi(t, z) = \{z^\ell + \sum_{j < \ell} \alpha_j z^j\} e^{\sum t_i z^i}. \quad (1.6)$$

The fact that M is a free $R[\partial, \partial^{-1}]$ -module, permits you to show that each oscillating function of type z^ℓ that satisfies (1.5) gives you a solution of (1.3). This function is then called a *wavefunction* of the KP-hierarchy. Notice that we can write $\psi(z) = P(\partial) e^{\sum t_i z^i}$. It is not difficult to see that the equations (1.5) for $L = P\partial P^{-1}$ are equivalent to the Sato–Wilson equations for P :

$$\partial_n(P)P^{-1} = -(P\partial^n P^{-1})_-. \quad (1.7)$$

Besides the space of oscillating functions M it is also convenient to have at one's disposal its “adjoint” space M^* consisting of all formal products

$$\left\{ \sum_{j \leq N} a_j z^j \right\} e^{-\sum t_i z^i}, \quad \text{with } a_j \in R \text{ for all } j.$$

The ring $R[\partial, \partial^{-1}]$ acts as expected on M^* by the natural extension of the actions

$$\begin{aligned} b\left\{ \left(\sum_j a_j z^j \right) e^{-\sum t_i z^i} \right\} &= \left(\sum_j b a_j z^j \right) e^{-\sum t_i z^i} \\ \partial\left\{ \left(\sum_j a_j z^j \right) e^{-\sum t_i z^i} \right\} &= \left(\sum_j \partial(a_j) z^j - \sum_j a_j z^{j+1} \right) e^{-\sum t_i z^i}. \end{aligned}$$

This is also a free $R[\partial, \partial^{-1}]$ -module, since we have

$$\left(\sum p_j (-\partial)^j \right) e^{-\sum t_i z^i} = \left(\sum p_j z^j \right) e^{-\sum t_i z^i}.$$

There is a bilinear pairing $\Re : M \times M^* \rightarrow R$ defined as follows: if $\phi = \phi(t, z) = \left(\sum_j a_j(t) z^j \right) e^{\sum t_i z^i}$ belongs to M and $\psi = \psi(t, z) = \left(\sum_k b_k(t) z^k \right) e^{-\sum t_i z^i}$ is an element of M^* , then we put

$$\begin{aligned} \Re(\phi, \psi) &:= \text{Res}_{z=0} \left\{ \left(\sum_j a_j(t) z^j \right) \left(\sum_k b_k(t) z^k \right) \right\} \\ &= \sum_{k \in \mathbb{Z}} a_{-k-1}(t) b_k(t), \end{aligned}$$

which is clearly a finite sum and hence belongs to R .

On $R[\partial, \partial^{-1}]$ we have an anti-algebra morphism called *taking the adjoint*. The adjoint of $P = \sum p_i \partial^i$ is given by

$$\begin{aligned} P^* &= \sum_i (-\partial)^i p_i = \sum_i (-)^i \sum_{k=0}^{\infty} \binom{i}{k} \partial^k (p_i) \partial^{i-k} \\ &= \sum_{\ell} \left\{ \sum_{k=0}^{\infty} (-)^{\ell+k} \binom{\ell+k}{k} \partial^k (p_{\ell+k}) \right\} \partial^{\ell}. \end{aligned}$$

There is an important connection between the bilinear form $\Re$ and taking the adjoint.

Theorem 1.1 *Let $\phi(t, z) = P(t, \partial) e^{\sum t_i z^i} \in M$ and $\psi(t, z) = Q(t, \partial) e^{-\sum t_i z^i} \in M^*$, then*

$$\begin{aligned} (P(t, \partial) Q(t, \partial)^*)_- &= \Re(\phi(t, z), \partial^{-1} \circ \psi(t, z)) \\ &:= - \sum_{n=0}^{\infty} \Re\left(\phi(t, z), \frac{\partial^n \psi(t, z)}{\partial x^n}\right) (-\partial)^{-n-1}. \end{aligned} \tag{1.8}$$

Proof. The proof consists of showing, by a direct computation, that the coefficients of ∂^{-n-1} on both sides of (1.8) are equal. Let $P = \sum_j a_j \partial^j$ and let $Q = \sum_k b_k \partial^k$. First we compute the coefficient $\Re(\phi(t, z), \frac{\partial^n \psi(t, z)}{\partial x^n})$. Since

$$\begin{aligned} \partial^n(\psi) &= \left\{ \sum_{\ell=0}^n \binom{n}{\ell} \sum_k \partial^{n-\ell}(b_k) (-z)^{k+\ell} \right\} e^{-\sum t_i z^i} \\ &= \left\{ \sum_p \left(\sum_{\ell=0}^n \binom{n}{\ell} (-)^p \partial^{n-\ell}(b_{p-\ell}) \right) z^p \right\} e^{-\sum t_i z^i}, \end{aligned}$$

we get that

$$\Re(\phi(t, z), \frac{\partial^n \psi(t, z)}{\partial x^n}) = \sum_p a_{-p-1} \sum_{\ell=0}^n \binom{n}{\ell} (-)^p \partial^{n-\ell}(b_{p-\ell}).$$

On the other hand we have

$$\begin{aligned} PQ^* &= \sum_{j,k} a_j \partial^j (-\partial)^k b_k = \sum_{j,k} (-)^k a_j \partial^{j+k} b_k \\ &= \sum_{j,k} (-)^k a_j \sum_{r=0}^{\infty} \binom{j+k}{r} \partial^r (b_k) \partial^{j+k-r} \\ &= \sum_s \left\{ \sum_j \sum_{r=0}^{\infty} \binom{r+s}{r} (-)^{r+s+j} a_j \partial^r (b_{r+s-j}) \right\} \partial^s. \end{aligned}$$

Hence the coefficient of ∂^{-n-1} , $n \geq 0$, equals

$$\begin{aligned} (-)^{n+1} \sum_j \sum_{r=0}^{\infty} \binom{r-n-1}{r} (-)^{r+j} a_j \partial^r (b_{r-j-n-1}) &= \\ &= (-)^{n+1} \sum_j \sum_{r=0}^n \binom{n}{r} (-)^j a_j \partial^r (b_{r-j-n-1}) \\ &= (-)^{n+1} \sum_j (-)^j a_j \sum_{\ell=0}^n \binom{n}{\ell} \partial^{n-\ell}(b_{-\ell-j-1}) \\ &= (-)^n \sum_p (-)^p a_{-p-1} \sum_{\ell=0}^n \binom{n}{\ell} \partial^{n-\ell}(b_{p-\ell}) \\ &= (-)^n \Re(\phi(t, z), \frac{\partial^n \psi(t, z)}{\partial x^n}). \end{aligned}$$

This proves the claim in the theorem. $\square$

If $\phi = P(t, \partial) e^{\sum t_i z^i}$ is an oscillating function of type z^ℓ , then we call $\phi^* = P(t, \partial)^* e^{-\sum t_i z^i}$ in M^* the adjoint of ϕ . It is a dual oscillating function of type $z^{-\ell}$. The foregoing theorem shows that for all $n \geq 0$

$$\Re(\phi(t, z), \frac{\partial^n \phi^*(t, z)}{\partial t_1^n}) = 0. \quad (1.9)$$

This property even characterizes ϕ^* among the dual oscillating functions of type $z^{-\ell}$. For, if $\psi = Q(t, \partial) e^{-\sum t_i z^i} \in M^*$ is such a function satisfying (1.9) with ϕ^* replaced by ψ , then we have according to the theorem

$$PQ^* = \partial^0 + (PQ^*)_- = 1.$$

In other words, $Q = (P^*)^{-1}$ and $\psi = \phi^*$. We will use this criterion later on and therefore we resume it in a

Lemma 1.1 *Let ϕ be an oscillating function of type z^ℓ and ψ a dual oscillating function of type $z^{-\ell}$. Then ψ is the adjoint of ϕ if and only if it satisfies*

$$\Re(\phi(t, z), \frac{\partial^n \psi(t, z)}{\partial t_1^n}) = 0 \quad \text{for all } n \geq 0.$$

If $\psi \in M$ is a wavefunction for the KP hierarchy, then its adjoint ψ^* satisfies a similar set of linear equations, viz.,

$$L^* \psi^* = z \psi^* \quad \text{and} \quad \partial_n(\psi^*) = -(L^n)_+(\psi^*), \quad (1.10)$$

where $L^* = (P \partial P^{-1})^* = (P^*)^{-1}(-\partial)P^*$. The first of these equations is a direct consequence of the definition for

$$(P^*)^{-1}(-\partial)P^*(P^*)^{-1}e^{-\sum t_i z^i} = (P^*)^{-1}(-\partial)e^{-\sum t_i z^i} = z(P^*)^{-1}e^{-\sum t_i z^i} = z\psi^*.$$

The second follows from the Sato–Wilson equations for P . By taking the adjoint of relation (1.7) we get

$$(\partial_n(P)P^{-1})^* = P^{*-1}\partial_n(P^*) = -(P\partial^n P^{-1})_-^* = -(P^{*-1}(-\partial)^n P^*)_- = -((L^*)^n)_-.$$

Since $\partial_n(P^{*-1}) = -P^{*-1}\partial^n(P^*)P^{*-1}$, these equations combine to give

$$\partial_n(P^{*-1})P^* = ((L^*)^n)_-. \quad (1.11)$$

As we have that

$$\begin{aligned} \partial_n(\psi^*) &= \partial_n(P^{*-1}e^{-\sum t_i z^i}) = \partial_n(P^{*-1})e^{-\sum t_i z^i} - z^n P^{*-1}e^{-\sum t_i z^i} \\ &= ((L^*)^n)_-\psi^* - (L^*)^n\psi^* = -((L^*)^n)_+\psi^*, \end{aligned}$$

which is exactly the second equation of (1.10). Reversely, if the adjoint of an oscillating function of type z^ℓ satisfies the equations in (1.10), then ψ is a wavefunction of the KP hierarchy. The equations in (1.10) namely imply equation (1.11) and by taking the adjoint of it, we get the Sato–Wilson equations for P .

In the present context, where the splitting of the oscillating functions in the exponential factor $e^{\sum t_i z^i}$ and a factor that has at most a pole at infinity is a formal affair. One also does not have at one's disposition the famous bilinear identity (see [9]):

$$Res_{z=0}\psi(t, z)\psi^*(s, z) = 0, \quad (1.12)$$

to characterize the oscillating functions ψ of type z^ℓ that are wavefunctions for the KP hierarchy. We will see in the next section that the relation (1.12) makes sense in the convergent setting of Segal and Wilson.

Another item in the theory of the KP hierarchy is the tau-function $\tau(t) := \tau((t_i))$. It was introduced by the Kyoto school, see [9], and relates to the wavefunction ψ and its adjoint wavefunction ψ^* by the formula's

$$\psi(t, z) = z^\ell \frac{\tau(t - [z^{-1}])}{\tau(t)} e^{\sum t_i z^i} \quad \text{and} \quad \psi^*(t, z) = z^{-\ell} \frac{\tau(t + [z^{-1}])}{\tau(t)} e^{-\sum t_i z^i}, \quad (1.13)$$

where $[\lambda] = (\lambda, \frac{1}{2}\lambda^2, \frac{1}{3}\lambda^3, \frac{1}{4}\lambda^4, \dots)$. In the general set-up of the oscillating functions its existence is by no means clear, but in the Segal–Wilson context such a τ exists. In order that the right-hand sides of the equations (1.13) are in M resp. M^* , there should hold the Taylor formula for τ . By applying this to (1.13) we get for all first order derivatives of τ

$$\frac{\partial_k(\tau)}{\tau} = Res_{z=0} z^k \left(\left(\frac{\partial}{\partial z} - \sum_{j=1}^{k-1} z^{-j-1} \partial_j \right) \log(z^{-\ell} e^{-\sum t_i z^i} \psi(t, z)) \right). \quad (1.14)$$

From this formula we see that if there are 2 tau-functions satisfying (1.13), then their quotient is independent of the variables (t_i) . If a wavefunction ψ of the KP hierarchy possesses a function τ satisfying (1.13) and if the bilinear form (1.12) makes sense, then this gives the following bilinear identity for the KP tau-functions:

$$Res_{z=0} \tau(t - [z^{-1}]) e^{\sum (t_i - s_i) z^i} \tau(s + [z^{-1}]) = 0. \quad (1.15)$$

2 The Segal–Wilson Grassmannian

An analytic approach to construct wavefunctions of the KP-hierarchy is considered in [22]. Segal and Wilson consider the Hilbert space

$$H = \left\{ \sum_{n \in \mathbb{Z}} a_n z^n \mid a_n \in \mathbb{C}, \sum_{n \in \mathbb{Z}} |a_n|^2 < \infty \right\},$$

with decomposition $H = H_+ \oplus H_-$, where

$$H_+ = \left\{ \sum_{n \geq 0} a_n z^n \in H \right\} \quad \text{and} \quad H_- = \left\{ \sum_{n < 0} a_n z^n \in H \right\}$$

and inner product $\langle \cdot | \cdot \rangle$ given by

$$\left\langle \sum_{n \in \mathbb{Z}} a_n z^n \mid \sum_{m \in \mathbb{Z}} b_m z^m \right\rangle = \sum_{n \in \mathbb{Z}} a_n \overline{b_n}.$$

To this decomposition is associated the Grassmannian $Gr(H)$ consisting of all closed subspaces W of H such that the orthogonal projection $p_+ : W \rightarrow H_+$ is Fredholm and the orthogonal projection $p_- : W \rightarrow H_-$ is Hilbert-Schmidt. The connected components of $Gr(H)$ are given by

$$Gr^{(\ell)}(H) = \left\{ W \in Gr(H) \mid p_+ : z^{-\ell} W \rightarrow H_+ \text{ has index zero} \right\}.$$

Each of these components is a homogeneous space for the group $Gl_{res}^{(0)}(H)$ of all bounded invertible operators $g : H \rightarrow H$ that decompose with respect to $H = H_+ \oplus H_-$ as

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

where a and d are Fredholm operators and b and c are Hilbert–Schmidt. For each $N > 1$, we consider the multiplicative group

$$\Gamma_+(N) = \left\{ \exp\left(\sum_{i \geq 1} t_i z^i\right) \mid t_i \in \mathbb{C}, \sum |t_i| N^i < \infty \right\}.$$

equipped with the uniform norm. These groups are nested in a natural way and the inductive limit is denoted by Γ_+ . The groups $\Gamma_+(N)$ act by multiplication on H and this gives a continuous injection of $\Gamma_+(N)$ into $Gl_{res}^{(0)}(H)$. The commuting flows from $\Gamma_+(N)$ lead to wavefunctions of the KP hierarchy for which the product in (1.6) is real and not formal.

Now we take for R the ring of meromorphic functions on $\Gamma_+(N)$ and for ∂_n the partial derivative w.r.t. the parameter t_n of $\Gamma_+(N)$. For each $W \in Gr^{(\ell)}(H)$, let $\Gamma_+^W(N)$ be given by

$$\Gamma_+^W(N) = \left\{ \gamma(t) = \exp\left(\sum_{i=1}^{\infty} t_i z^i\right) \in \Gamma_+(N) \mid p_+ : \gamma^{-1} z^{-\ell} W \rightarrow H_+ \text{ is a bijection} \right\}.$$

In a similar way as in [22], one shows that $\Gamma_+^W(N)$ is nonempty. For $\gamma \in \Gamma_+^W(N)$, let $P_W(t, z)$ be z^ℓ times the inverse image of 1 under the projection $p_+ : \gamma^{-1}z^{-\ell}W \rightarrow H_+$. Then we associate to W an oscillating function $\psi_W(t, z) = P_W(t, z)e^{\sum_{i=1}^\infty t_i z^i}$ of type z^ℓ , which has the properties that it is defined on a dense open subset of $\Gamma_+(N)$ and that its boundary value at $|z| = 1$ belongs to W . Moreover, it is known that the range of ψ_W spans a dense subspace of W . A crucial property of the elements $\gamma(t)$ in this dense open subset of $\Gamma_+(N)$ is that

$$W \cap (z^\ell H_+)^{\perp} \gamma(t) = \{0\}.$$

By exploiting this property, one shows that ψ_W is a wavefunction for the KP hierarchy. Hence, if we write $\psi_W(t, z) = P_W(t, \partial) \cdot e^{\sum t_i z^i}$ with $P_W \in R[\partial, \partial^{-1}]$, then $L_W = P_W \partial P_W^{-1}$ is a solution of the KP-hierarchy. Each component of $Gr(H)$ generates in this way the same set of solutions of the KP-hierarchy, so it would suffice, as is done in [22], to consider only $Gr^{(0)}(H)$. However, we need to consider here all components.

Let $W \in Gr^{(\ell)}(H)$, then $W^\perp$ is a closed subspace of H , moreover $p_- : W^\perp \rightarrow H_-$ is a Fredholm operator of index $-\ell$ and $p_+ : W^\perp \rightarrow H_+$ is a Hilbert–Schmidt operator. Interchanging the role of H_+ and H_- , we see that $W^\perp$ is a plane in the adjoint Grassmannian $Gr^*(H)$ consisting of planes U for which p_- is a Fredholm operator and p_+ is a Hilbert–Schmidt operator. The connected components of $Gr^*(H)$ are also homogeneous spaces of the group $Gl_{res}^{(0)}(H)$ introduced above. On $Gr^*(H)$ we consider however the commuting flows that are the adjoint of the ones on $Gr(H)$. Namely, for each $N > 1$, we take the group

$$\Gamma_-(N) = \{\exp(\sum_{i \geq 1} r_i z^{-i}) \mid r_i \in \mathbb{C}, \sum |r_i| N^i < \infty\}.$$

and as the analogue of $\Gamma_+^W(N)$ we take here

$$\Gamma_-^{W^\perp}(N) = \{g(r) = \exp(\sum_{i=1}^\infty r_i z^{-i}) \in \Gamma_-(N) \mid p_- : g^{-1}z^\ell W^\perp \rightarrow H_- \text{ is a bijection}\}.$$

One verifies directly that

$$\Gamma_-^{W^\perp}(N) = \{(\gamma^*)^{-1} \mid \gamma \in \Gamma_+^W(N)\}.$$

In particular it is a non-empty open dense subset of $\Gamma_-(N)$. Now we can repeat the construction of the wavefunction ψ_W with $W \in Gr(H)$ replaced by $W^\perp \in Gr^*(H)$ and $\Gamma_+(N)$ by $\Gamma_-(N)$, thus we obtain a function $\psi_{W^\perp}$ of the form

$$\begin{aligned} \psi_{W^\perp}(r, z) &= \psi_{W^\perp}(g(r), z) = \left\{ \sum_{s \geq \ell-1} \alpha_s(r) z^s \right\} g(r) \\ &= \hat{\psi}_{W^\perp}(g(r), z) g(r). \end{aligned}$$

Here $\alpha_{\ell-1} = 1$ and $g(r)$ belongs to $\Gamma_-^{W^\perp}(N)$. For all such $g(r)$ the boundary value of $\psi_{W^\perp}$ on $|z| = 1$ belongs to $W^\perp$. This last property also holds for

$$\begin{aligned} \left(\frac{\partial}{\partial r_1} \right)^n \psi_{W^\perp}(g(r), z) &= \left\{ \sum_{s \geq \ell-1} \sum_{p=0}^n \binom{n}{p} \frac{\partial^{n-p} \alpha_s(r)}{\partial r_1^{n-p}} z^{s-p} \right\} g(r) \\ &:= \left\{ \sum_{s \geq \ell-n-1} \alpha_s^{(n)}(r) z^s \right\} g(r). \end{aligned}$$

Since the boundary value on $|z| = 1$ belongs to W , we get for all $\gamma(t) \in \Gamma_+^W(N)$ and all $g(r) \in \Gamma_-^{W^\perp}(N)$ the equality

$$\langle \psi_W(\gamma(t), z) | \left(\frac{\partial}{\partial r_1} \right)^n \psi_{W^\perp}(g(r), z) \rangle = 0. \quad (2.1)$$

Recall that the innerproduct is also equal to

$$\langle f(z) | h(z) \rangle = \frac{1}{2\pi i} \int_{|z|=1} f(z) \overline{h(z)} \frac{dz}{z}. \quad (2.2)$$

For each $g(r) \in \Gamma_-^{W^\perp}(N)$ there is a $\gamma(s) \in \Gamma_+^W(N)$ such that $g(r) = \gamma(-\bar{s})$. Hence we get on $|z| = 1$

$$\begin{aligned} \overline{\left(\frac{\partial}{\partial r_1} \right)^n \psi_{W^\perp}(g(r), z)} &= \left\{ \sum_{p \geq \ell - n - 1} \overline{\alpha_p^{(n)}(r) z^{-p}} \right\} e^{\sum_{i=1}^{\infty} \bar{r}_i z^i} \\ &= \left\{ \sum_{j \leq n - \ell + 1} \overline{\alpha_{-j}^{(n)}(r) z^j} \right\} e^{\sum_{i=1}^{\infty} \bar{r}_i z^i} \\ &= \left\{ \sum_{j \leq n - \ell + 1} \overline{\alpha_{-j}^{(n)}(-\bar{s}) z^j} \right\} e^{\sum_{i=1}^{\infty} s_i z^i} \\ &= (-)^n \left(\frac{\partial}{\partial s_1} \right)^n \left\{ \sum_{j \leq -\ell + 1} \overline{\alpha_j(-\bar{s}) z^j} \gamma(s) \right\}. \end{aligned}$$

In other words $\overline{\psi_{W^\perp}(\gamma(-\bar{s}), z)}$ is a dual oscillating function on M^* of type $z^{1-\ell}$. If we substitute this into formula (2.2) and we put $s_k = t_k$ for all $k \geq 1$, then we obtain from (2.1) for all $n \geq 0$

$$\Re(\psi_W(\gamma(t), z), \partial^n(z^{-1} \overline{\psi_{W^\perp}(\gamma(-\bar{t}), z)})) = 0.$$

So $z^{-1} \overline{\psi_{W^\perp}(\gamma(-\bar{t}), z)}$ satisfies the requirements in Lemma 1.1 and we can say

Theorem 2.1 *The adjoint wavefunction of $\psi_W(\gamma(t), z)$ is $z^{-1} \overline{\psi_{W^\perp}(\gamma(-\bar{t}), z)}$.*

A consequence of this theorem is that the adjoint wavefunction of a wavefunction of the Segal–Wilson class, falls again in this class. This is by no means obvious from the defining formula. Moreover, we can give a meaning to the bilinear formula for the Segal–Wilson wavefunctions, viz., for all relevant t and s one has

$$\langle \psi_W(t, z) | \overline{z \psi_W^*(s, z)} \rangle = 0. \quad (2.3)$$

We can use the fact that the boundary values of $\psi_W(t, z)$ on $|z| = 1$ are lying dense in W and that the boundary values of $\psi_W(s, z)$ at $|z| = 1$ also belong to H to characterize inclusions inside $Gr(H)$. This gives you so-called modified equations. There holds namely

Proposition 2.1 *Let V, W be planes in $Gr(H)$. Then $V \subset W$ (and hence $W^\perp \subset V^\perp$) if and only if*

$$\langle \psi_V(t, z) | \overline{z \psi_W^*(s, z)} \rangle = 0.$$

Next we recall how the tau-functions arise in the Segal–Wilson setting. This starts with a convenient description of components $Gr^{(\ell)}(H)$ of the Grassmannian. Let $\mathfrak{P}_\ell$ be the collection of embeddings $w : z^\ell H_+ \rightarrow H$ such that with respect to the decomposition $H = (z^\ell H_+) \oplus (z^\ell H_+)^\perp$ the operator w has the form $w = \begin{pmatrix} w_+ \\ w_- \end{pmatrix}$, with w_- a Hilbert–Schmidt operator and $w_+ - \text{Id}$ a trace class operator. Then $\mathfrak{P}_\ell$ is a natural way a fibre bundle over $Gr^{(\ell)}(H)$ with fiber the group

$$\mathfrak{T}_\ell = \{t \in \text{Aut}(z^\ell H_+) | t - \text{Id} \text{ of trace class}\}.$$

To lift the action of $GL_{res}^{(0)}(H)$ on $Gr^{(\ell)}(H)$ to one on $\mathfrak{P}_\ell$, one has to pass to an extension Gl of $Gr^{(\ell)}(H)$. It is defined by

$$Gl = \{(g, q) | g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, g \in GL_{res}^{(0)}(H), q \in \text{Aut}(z^\ell H_+), aq^{-1} - \text{Id} \text{ is trace class}\}.$$

This group acts by $w \mapsto gwq^{-1}$ on $\mathfrak{P}_\ell$. The elements of both $\Gamma_+(N)$ and $\Gamma_-(N)$ embed in a natural way into Gl through

$$\gamma_+ = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \mapsto (\gamma_+, a) \text{ and } \gamma_- = \begin{pmatrix} p & 0 \\ r & s \end{pmatrix} \mapsto (\gamma_-, p).$$

This we assume through this paper. For each $w \in \mathfrak{P}_\ell$, we define $\tau_w : Gl \rightarrow \mathbb{C}$ by

$$\tau_w((g, q)) = \det((g^{-1}wq)_+).$$

If t belongs to $\mathfrak{T}_\ell$, then there holds $\tau_{w \circ t} = \det(t)\tau_w$. The restriction of τ_w to $\Gamma_+(N)$ gives you the required tau-function. For, if $q_\zeta = 1 - \frac{\zeta}{\zeta} \in \Gamma_+(N)$, then there holds

Theorem 2.2 *Let W be a plane in $Gr^{(\ell)}(H)$ and let $\psi_W(t, z) = P_W(t, z)e^{\sum t_i z^i}$ be the corresponding wavefunction of type z^ℓ . Then we have*

$$\begin{aligned} (a) \quad & P_W(t, \zeta) = \zeta^\ell \frac{\tau_w(\gamma(t)q_\zeta)}{\tau_w(\gamma(t))} = \zeta^\ell \frac{\tau_W(t - [\zeta^{-1}])}{\tau_W(t)} \quad \text{and} \\ (b) \quad & \text{the adjoint wavefunction } \psi_W^* \text{ satisfies} \\ & \psi_W^*(t, \zeta) = \zeta^{-\ell} \frac{\tau_w(\gamma(t)q_\zeta^{-1})}{\tau_w(\gamma(t))} e^{-\sum_{i=1}^\infty t_i \zeta^i} = \zeta^{-\ell} \frac{\tau_W(t + [\zeta^{-1}])}{\tau_W(t)} e^{-\sum_{i=1}^\infty t_i \zeta^i}. \end{aligned}$$

Property (a) was shown in [22] and the relation in (b) is the combination of theorem 1.1 and the computations in [10]. This theorem shows that in the Segal–Wilson setting we have the equations in (1.13).

3 Elementary Bäcklund–Darboux Transformations

In this section we want to consider a certain type of Bäcklund–Darboux transformations the so-called elementary ones. For some literature on Bäcklund–Darboux transformations we refer the reader to [2], [5], [15], [17] and [18].

Let ψ be a wavefunction of type z^ℓ , not necessarily in the Segal–Wilson setting. Like in [17] and [18], we define elementary Bäcklund–Darboux transformations that produce new wavefunctions of type $z^{\ell \pm 1}$. For $q, r \in R$, one considers the first order differential operators $q\partial q^{-1}$ and $r^{-1}\partial r$. An obvious question now is: When are $q\partial q^{-1}\psi$ and $r^{-1}\partial^{-1}r\psi$

again wavefunctions? The answer is given by the following theorem. But first notice that $r^{-1}\partial^{-1}r\psi$ is well and uniquely defined. If $\psi = P(z)e^{\sum t_i z^i} = P(\partial)e^{\sum t_i z^i}$, then

$$\begin{aligned} r^{-1}\partial^{-1}r\psi &= r^{-1}\partial^{-1}rP(\partial)e^{\sum t_i z^i} \\ &= r^{-1}\sum_{k=0}^{\infty}(-)^k \frac{\partial^k(rP(\partial))}{\partial x^k} \partial^{-k-1} e^{\sum t_i z^i} \\ &= r^{-1}\sum_{k=0}^{\infty}(-)^k \frac{\partial^k(rP(z))}{\partial x^k} z^{-k-1} e^{\sum t_i z^i}. \end{aligned}$$

Theorem 3.1 *Let ψ be a wavefunction of type z^ℓ and let ψ^* be its adjoint wave function. Then the following holds:*

(a) *Let q and $r \in R$ satisfy*

$$\partial_n(q) = (L^n)_+(q), \quad (3.1)$$

$$\partial_n(r) = -(L^n)^*_+(r) \quad \text{for all } n = 1, 2, \dots \quad (3.2)$$

Then $q\partial q^{-1}\psi$ (resp. $r^{-1}\partial^{-1}r\psi$) is a wavefunction of type $z^{\ell+1}$ (resp. $z^{\ell-1}$) and $-q^{-1}\partial^{-1}q\psi^$ (resp. $-r\partial r^{-1}\psi^*$) is its adjoint wavefunction.*

(b) *If both q_1 and q_2 (resp. r_1 and r_2) satisfy (3.1) (resp. (3.2)) such that*

$$q_1\partial q_1^{-1} = q_2\partial q_2^{-1} \quad (\text{resp. } r_1^{-1}\partial r_1 = r_2^{-1}\partial r_2),$$

then $q_1 = \lambda q_2$ (resp. $r_1 = \lambda r_2$) for certain $\lambda \in \mathbb{C}$.

We say that q (resp. r) is an *eigenfunction* (resp. *adjoint eigenfunction*) of the Lax operator L if it satisfies (3.1) (resp. (3.2)).

Recall that there is a unique pseudodifferential operator P such that $\psi = P(e^{\sum t_i z^i})$. It has the form

$$P = \partial^\ell + \sum_{j < \ell} p_j \partial^j = \{1 + \sum_{s < 0} p_{\ell+s} \partial^s\} \partial^\ell, \quad (3.3)$$

which satisfies the Sato–Wilson equation (1.7). Let τ_q (resp. τ_r) be the tau-function corresponding to $q\partial q^{-1}\psi$ (resp. $r^{-1}\partial^{-1}r\psi$), by using formula (1.13) one easily deduces that

$$\frac{\partial \log q}{\partial x} = \frac{\partial \log \tau_q / \tau}{\partial x} \quad (\text{resp. } \frac{\partial \log r}{\partial x} = \frac{\partial \log \tau_r / \tau}{\partial x}). \quad (3.4)$$

Next we consider the operators P_q and P_r defined by

$$P_q := q\partial q^{-1}P \quad \text{and} \quad P_r := r^{-1}\partial^{-1}rP. \quad (3.5)$$

To Prove part (a) of Theorem 3.1, one has to show that the P_q and the P_r also satisfy the Sato–Wilson equations. To do so, we need some properties of the ring $R[\partial, \partial^{-1}]$ of pseudodifferential operators with coefficients from R . We resume them in a lemma

Lemma 3.1 *If f belongs to R and Q to $R[\partial, \partial^{-1}]$, then the following identities hold*

$$(a) \quad (Qf)_- = Q_-f,$$

$$(b) \quad (fQ)_- = fQ_-,$$

$$(c) \quad \text{Res}_\partial(Qf) = \text{Res}_\partial(fQ) = f \text{Res}_\partial(Q),$$

- (d) $(\partial Q)_- = \partial Q_- - \text{Res}_\partial(Q),$
- (e) $(Q\partial)_- = Q_- \partial - \text{Res}_\partial(Q),$
- (f) $(Q\partial^{-1})_- = Q_- \partial^{-1} + \text{Res}_\partial(Q\partial^{-1})\partial^{-1},$
- (g) $(\partial^{-1}Q)_- = \partial^{-1}Q_- + \partial^{-1} \text{Res}_\partial(Q^* \partial^{-1}),$
- (h) $\text{Res}_\partial(Qf\partial^{-1}) = Q_+(f),$
- (i) $\text{Res}_\partial(\partial^{-1}fQ) = Q_+^*(f).$

Since the proof of this lemma consists of straightforward calculations, we leave this to the reader. Now we can show

Proposition 3.1 *The operators P_q and P_r satisfy the Sato-Wilson equations.*

Proof. If we denote $\frac{\partial}{\partial t_n}$ by ∂_n , then we get for $P_q = q\partial q^{-1}P$ that

$$\begin{aligned} \partial_n(P_q)P_q^{-1} &= \partial_n(q\partial q^{-1})q\partial^{-1}q^{-1} + q\partial q^{-1}\partial_n(P)P^{-1}q\partial^{-1}q^{-1} \\ &= -q\partial q^{-1}(L^n)_-q\partial^{-1}q^{-1} + \partial_n(q\partial q^{-1})q\partial^{-1}q^{-1}. \end{aligned}$$

Now we apply successively the identities from Lemma 3.1 to the first operator of the right-hand side

$$\begin{aligned} q\partial q^{-1}(L^n)_-q\partial^{-1}q^{-1} &= q\partial(q^{-1}L^nq)_-\partial^{-1}q^{-1} &= \\ q\partial(q^{-1}L^nq\partial^{-1})_-q^{-1} &- q\partial\text{Res}_\partial(q^{-1}L^nq\partial^{-1})\partial^{-1}q^{-1} &= \\ q(\partial q^{-1}L^nq\partial^{-1})_-q^{-1} &+ q\text{Res}_\partial(q^{-1}L^nq\partial^{-1})q^{-1} &- \\ q\partial\text{Res}_\partial(q^{-1}L^nq\partial^{-1})\partial^{-1}q^{-1} &= (q\partial q^{-1}L^nq\partial^{-1}q^{-1})_- &+ \\ q^{-1}\text{Res}_\partial(L^nq\partial^{-1}) &- q\partial q^{-1}\text{Res}_\partial(L^nq\partial^{-1})\partial^{-1}q^{-1} &- \\ (q\partial q^{-1}L^nq\partial^{-1}q^{-1})_- &+ (L^n)_+(q)q^{-1} - q\partial q^{-1}(L^n)_+(q)\partial^{-1}q^{-1}. \end{aligned}$$

On the other hand

$$\partial_n(q\partial q^{-1})q\partial^{-1}q^{-1} = \partial_n(q)q^{-1} - q\partial q^{-2}\partial_n(q)q\partial^{-1}q^{-1}.$$

Thus we see that, if $\partial_n(q) = (L^n)_+(q)$, the operator P_q satisfies the Sato-Wilson equation

$$\partial_n(P_q)P_q^{-1} = -(P_q\partial^n P_q^{-1})_-. \quad (3.6)$$

For P_r , we proceed in a similar fashion

$$\begin{aligned} \partial_n(P_r)P_r^{-1} &= -r^{-1}\partial^{-1}r(L^n)_-r\partial r + \partial_n(r^{-1}\partial^{-1}r)r^{-1}\partial r \\ &= -r^{-1}\partial^{-1}(rL^n r^{-1})_- \partial r + -\partial_n(r)r^{-1} + r^{-1}\partial^{-1}(\partial_n(r)r^{-1})\partial r. \end{aligned}$$

Now we successively apply Lemma 3.1 (g) and (c) and (1.7) to the first term of the right hand side of this equation

$$\begin{aligned} -r^{-1}\partial^{-1}(rL^n r^{-1})_- \partial r &= -r^{-1}\{(\partial^{-1}rL^n r^{-1})_- - \partial^{-1}\text{Res}_\partial(r^{-1}(L^n)_+^* r\partial^{-1})\}\partial r \\ &= -r^{-1}(\partial^{-1}rL^n r^{-1})_- \partial r + r^{-1}\partial^{-1}r^{-1}(L^n)_+^*(r)\partial r \\ &= -r^{-1}\{(\partial^{-1}rL^n r^{-1}\partial)_- + \text{Res}_\partial(\partial^{-1}rL^n r^{-1})\}r + r^{-1}\partial^{-1}r^{-1}(L^n)_+^*(r)\partial r \\ &= -(r^{-1}\partial^{-1}rL^n r^{-1}\partial)_- - r^{-1}(L^n)_+^*(r) + r^{-1}\partial^{-1}r^{-1}(L^n)_+^*(r)\partial r. \end{aligned}$$

Since $\partial_n(r) = -(L^n)^*(r)$, we see that the last two terms cancel $\partial_n(r^{-1}\partial r)r^{-1}\partial r$ and thus we have obtained the Sato–Wilson equation for P_r

$$\partial_n(P_r)P_r = -(P_r\partial^n P_r^{-1})_-. \quad (3.7)$$

This concludes the proof of Proposition 3.1. $\square$

Thus the proof of part (a) of Theorem 3.1 has been completed. To prove part (b), we first calculate

$$\partial \left(\frac{q_1}{q_2} \right) = \frac{q_1}{q_2} \frac{\partial(q_1)}{q_1} - \frac{q_1}{q_2} \frac{\partial(q_2)}{q_2} = 0$$

So $\partial^k(\frac{q_1}{q_2}) = 0$ and $\partial^k(q_1) = \partial^k(\frac{q_1}{q_2}q_2) = \frac{q_1}{q_2}\partial^k(q_2)$. Now let $B_\ell = (L^\ell)_+$, then

$$\begin{aligned} \partial_n \left(\frac{q_1}{q_2} \right) &= \frac{B_n(q_1)}{q_2} - \frac{q_1}{q_2^2} B_n(q_2) \\ &= \frac{1}{q_2} B_n \left(\frac{q_1}{q_2} q_2 \right) - \frac{q_1}{q_2^2} B_n(q_2) \\ &= \frac{q_1}{q_2^2} B_n(q_2) - \frac{q_1}{q_2^2} B_n(q_2) \\ &= 0 \end{aligned}$$

for all $n = 1, 2, \dots$. So we conclude that $q_1 = \lambda q_2$ for a certain $\lambda \in \mathbb{C}$. Uniqueness for r follows analogously. This finishes the proof of Theorem 3.1.

Suppose now $q\partial q^{-1}\psi$ and $r^{-1}\partial^{-1}r\psi$ are again wavefunctions, but q is not an eigenfunction and r is not an adjoint eigenfunction. Since P_q and P_r satisfy the Sato–Wilson equation,

$$\begin{aligned} \partial_n(q) - (L^n)_+(q) - q\partial q^{-1}\{\partial_n(q) - (L^n)_+(q)\}\partial^{-1} &= 0, \\ \partial_n(r) + (L^n)_+^*(r) - \partial\{\partial_n(r) + (L^n)_+^*(r)\}r^{-1}\partial r &= 0 \end{aligned}$$

and thus

$$\begin{aligned} \partial(q^{-1}\{\partial_n(q) - (L^n)_+(q)\}) &= 0, \\ \partial(r^{-1}\{\partial_n(r) + (L^n)_+^*(r)\}) &= 0. \end{aligned}$$

So

$$\begin{aligned} \partial_n(q) - (L^n)_+(q) &= f_n(t_2, t_3, \dots)q, \\ \partial_n(r) + (L^n)_+^*(r) &= g_n(t_2, t_3, \dots)r. \end{aligned} \quad (3.8)$$

Let again $B_\ell = (L^\ell)_+$, and write down the compatibility conditions for q , i.e., $\partial_n(\partial_\ell(q)) = \partial_\ell(\partial_n(q))$. One has

$$\begin{aligned} \partial_n(\partial_\ell(q)) &= \partial_n(B_\ell(q) + f_\ell q) \\ &= \partial_n(B_\ell)(q) + B_\ell(\partial_n(q)) + \partial_n(f_\ell)q + f_\ell\partial_n(q) \\ &= \partial_n(B_\ell)(q) + B_\ell B_n(q) + B_\ell(f_n q) + \partial_n(f_\ell)q + f_\ell B_n(q) + f_\ell f_n q \end{aligned}$$

and a similar relation with n and ℓ interchanged, so we find

$$\begin{aligned} (\partial_n(B_\ell) - \partial_\ell(B_n) + B_\ell B_n - B_n B_\ell)(q) + \\ B_\ell(f_n q) - B_n(f_\ell q) + f_\ell B_n(q) - f_n B_\ell(q) + \\ \partial_n(f_\ell)q - \partial_\ell(f_n)q = 0. \end{aligned}$$

Because the Zakharov–Shabat equations (1.4) hold for the B_ℓ 's and f_m is independent of $x = t_1$ one finds that

$$\partial_n(f_\ell) = \partial_\ell(f_n)$$

and hence if it is possible to integrate to all times t_j one can find a function $F(t)$ such that

$$f_n(t) = F(t)^{-1} \partial_n F(t). \quad (3.9)$$

Now choose

$$\hat{q} = F(t)^{-1} q,$$

then $\partial_n(\hat{q}) = B_n(\hat{q})$. A similar argument holds for r . So we have under certain conditions, viz. certain integrability of functions, shown that one can find an eigenfunction $\hat{q}$ (resp. adjoint eigenfunction $\hat{r}$) which produces the same elementary Bäcklund–Darboux transformation, i.e.

$$q \partial q^{-1} = \hat{q} \partial \hat{q}^{-1} \quad \text{resp.} \quad r \partial r^{-1} = \hat{r} \partial \hat{r}^{-1}.$$

We will show that in the Segal–Wilson case the above conditions are satisfied. In that case we will explicitly construct the (adjoint) eigenfunctions $\hat{q}$ and $\hat{r}$.

Clearly the Bäcklund–Darboux operators of Theorem 3.1 are invertible. Moreover, we prove the following useful lemma.

Lemma 3.2 *Let $q, r \in R$, then q (resp. r) satisfies (3.1) (resp. (3.2)) if and only if q^{-1} (resp. r^{-1}) satisfies*

$$\begin{aligned} \partial_n(q^{-1}) &= -(q \partial q^{-1} L^n q \partial^{-1} q^{-1})_+^*(q^{-1}), \\ \partial_n(r^{-1}) &= (r^{-1} \partial^{-1} r L^n r^{-1} \partial r)_+(r^{-1}), \quad \text{respectively.} \end{aligned}$$

Proof. Suppose (3.1) holds, then

$$\begin{aligned} \partial_n(q^{-1}) &= -q^{-2}(L^n)_+(q) \\ &= -q^{-2} \text{Res}_\partial(\partial^{-1} L^n)^* \\ &= -\text{Res}_\partial(q^{-1} \partial^{-1} q L^{n*} q^{-1}) \\ &= -\text{Res}_\partial(q^{-1} \partial^{-1} q L^{n*} q^{-1} \partial q q^{-1} \partial^{-1}) \\ &= -(q^{-1} \partial^{-1} q L^{n*} q^{-1} \partial q)_+(q^{-1}) \\ &= -(q \partial q^{-1} L^n q \partial^{-1} q^{-1})_+^*(q^{-1}). \end{aligned}$$

The second equation of this Lemma is proven in a similar fashion. $\square$

4 Trivial Bäcklund–Darboux Transformations

Besides the elementary Bäcklund–Darboux transformations one also has trivial Bäcklund–Darboux transformations. These transformations do not change the Lax operator L . They consist of multiplying a wavefunction with a function of the form

$$R(z) = z^k \left(1 + \sum_{j=1}^{\infty} r_j z^{-j}\right) \quad \text{with } r_j \in \mathbb{C}. \quad (4.1)$$

If $\psi(t, z) = P(t, \partial) e^{\sum_i t_i z^i}$ is a wavefunction of type z^ℓ , then

$$\begin{aligned}\psi_R(t, z) &= R(z) \psi(t, z) = P(t, \partial) R(\partial) e^{\sum_i t_i z^i} \\ &= z^k \left(1 + \sum_{j=1}^{\infty} r_j z^{-j} \right) \psi(t, z)\end{aligned}$$

is the new wavefunction of type $z^{k+\ell}$. Its adjoint wavefunction is of the form

$$\psi_R^*(t, z) = P(t, \partial)^{* -1} R(\partial)^{* -1} e^{-\sum_i t_i z^i} = R(z)^{-1} \psi^*(t, z).$$

Since all r_j 's are constants, this does not change the form of L , but it clearly changes the tau-function. Rewrite $R(z)$ as in (4.1) as follows

$$R(z) = z^k e^{-\sum_{j=1}^{\infty} \frac{q_j}{j} z^{-j}}, \quad (4.2)$$

then

$$\begin{aligned}\psi_R(t, z) &= z^k e^{-\sum_{j=1}^{\infty} \frac{q_j}{j} z^{-j}} \psi(t, z) \\ &= z^{k+\ell} \frac{\tau(t - [z^{-1}])}{\tau(t)} e^{-\sum_{j=1}^{\infty} \frac{q_j}{j} z^{-j}} e^{\sum_{i=1}^{\infty} t_i z^i} \\ &= z^{k+\ell} \frac{\tau(t - [z^{-1}]) e^{\sum_{j=1}^{\infty} q_j (t_j - \frac{z^{-j}}{j})}}{\tau(t) e^{\sum_{j=1}^{\infty} q_j t_j}} e^{\sum_{i=1}^{\infty} t_i z^i},\end{aligned}$$

which suggests that $\tau_R(t) := \tau(t) e^{\sum_{j=1}^{\infty} q_j t_j}$ is the to ψ_R corresponding tau-function. We will show this in the Segal-Wilson context. First of all, one notices that the multiplication by z^k does not change the tau-function. Hence we may assume that $k = 0$ and we consider the action of

$$\rho = \exp\left(-\sum_{j=1}^{\infty} \frac{q_j}{j} z^{-j}\right) \in \Gamma_-(N)$$

on $\mathfrak{P}_\ell$. The operator ρ decomposes with respect to $H = z^\ell H_+ \oplus (z^\ell H_+)^\perp$ as $\rho = \begin{pmatrix} \alpha & 0 \\ \beta & \delta \end{pmatrix}$.

Hence its action on $w = \begin{pmatrix} w_+ \\ w_- \end{pmatrix} \in \mathfrak{P}_\ell$ is

$$\rho \circ w \circ \alpha^{-1} = \begin{pmatrix} \alpha w_+ \alpha^{-1} \\ (\beta w_+ + \delta w_-) \alpha^{-1} \end{pmatrix}.$$

Let $\gamma^{-1} = \exp(-\sum_i t_i z^i) \in \Gamma_+(N)$ decompose with respect to $H = z^\ell H_+ \oplus (z^\ell H_+)^\perp$ as $\gamma^{-1} = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}$. Then the tau-function corresponding to ρW , where $W = \text{Im}(w)$, $w \in \mathfrak{P}_\ell$, is by definition

$$\tau_{\rho \circ w \circ \alpha^{-1}}(\gamma) = \det((\gamma^{-1} \rho w \alpha^{-1} a^{-1})_+).$$

Since γ^{-1} and ρ commute we have the relation

$$\alpha a = a \alpha + b \beta \quad \text{or equivalently} \quad \alpha a \alpha^{-1} a^{-1} = I + b \beta \alpha^{-1} a^{-1},$$

so that we see that the operator $\alpha a \alpha^{-1} a^{-1}$ is of the form “*identity + trace-class*” and hence has a determinant. Thus we get for the tau-function of ρW that

$$\begin{aligned}\tau_{\rho \circ w \circ \alpha^{-1}}(\gamma) &= \det((\gamma^{-1} \rho w a^{-1} \alpha^{-1})_+) \det(\alpha a \alpha^{-1} a^{-1}) \\ &= \det((\rho \gamma^{-1} w a^{-1} \alpha^{-1})_+) \det(\alpha a \alpha^{-1} a^{-1}). \\ &= \tau_w(\gamma) \det(\alpha a \alpha^{-1} a^{-1})\end{aligned}$$

Hence we merely have to show that

$$B(\alpha, a) = \det(\alpha a \alpha^{-1} a^{-1})$$

is the required function. This we will prove in a few steps. First one notices that if

$\rho_1 = \begin{pmatrix} \alpha_1 & 0 \\ \beta_1 & \delta_1 \end{pmatrix}$ and $\rho_2 = \begin{pmatrix} \alpha_2 & 0 \\ \beta_2 & \delta_2 \end{pmatrix}$ are in $\Gamma_-(N)$, then

$$\begin{aligned}B(\alpha_1 \alpha_2, a) &= \det(\alpha_1 \alpha_2 a \alpha_2^{-1} \alpha_1^{-1} a^{-1}) = \det(\alpha_2 a \alpha_2^{-1} a^{-1} a \alpha_1^{-1} a^{-1} \alpha_1) \\ &= \det(\alpha_2 a \alpha_2^{-1} a^{-1}) \det(a \alpha_1^{-1} a^{-1} \alpha_1) \\ &= B(\alpha_2, a) B(\alpha_1, a).\end{aligned}$$

Likewise one shows for $\gamma_i^{-1} = \begin{pmatrix} a_i & b_i \\ 0 & d_i \end{pmatrix} \in \Gamma_+(N)$ that

$$B(\alpha, a_1 a_2) = B(\alpha, a_1) B(\alpha, a_2).$$

Hence if we put $\rho_i = \exp(-\frac{q_i}{i} z^{-i}) = \begin{pmatrix} \alpha_i & 0 \\ \beta_i & \delta_i \end{pmatrix}$, then $\rho = \lim_{N \rightarrow \infty} \rho_1 \rho_2 \cdots \rho_N$. The form B is continuous in α , so that we get

$$B(\alpha, a) = \prod_{i=1}^{\infty} B(\alpha_i, a)$$

and thus we merely have to prove the formula for ρ_i . If

$$\gamma_i^{-1} = \exp(-t_i z^i) = \begin{pmatrix} a_i & b_i \\ 0 & d_i \end{pmatrix} \in \Gamma_+(N),$$

then $\gamma = \lim_{N \rightarrow \infty} \gamma_1 \gamma_2 \cdots \gamma_N$. As B is also continuous and multiplicative in a , we end up with the formula

$$B(\alpha, a) = \prod_{i,j \geq 1} B(\alpha_i, a_j).$$

Therefore we only have to prove the formula for $\rho = \rho_i$ and $\gamma = \gamma_j$. Clearly the map $q_i \rightarrow B(\alpha_i, a_j)$ defines a continuous morphism $\chi_i^{(j)}$ from $\mathbb{C}$ to $\mathbb{C}^\times$ and therefore it has the form

$$\chi_i^{(j)}(q_i) = e^{\beta_{ij} q_i},$$

for some $\beta_{ij} \in \mathbb{C}$, depending on γ_j . Likewise the map $t_j \rightarrow B(\alpha_i, a_j)$ defines a continuous morphism $\phi_j^{(i)}$ from $\mathbb{C}$ to $\mathbb{C}^\times$ that can be written as

$$\phi_j^{(i)}(t_j) = e^{\alpha_{ij} t_j},$$

for some $\alpha_{ij} \in \mathbb{C}$, depending on ρ_j . Hence we can say that for all $i \geq 1$ and $j \geq 1$

$$B(\alpha_i, a_j) = e^{a_{ij} t_i q_j} \quad \text{with } a_{ij} \in \mathbb{C}.$$

The $\{a_{ij}\}$ can easily be determined by computing the lowest non-trivial term in $B(\alpha_i, a_j)$. Since

$$b_j = \begin{pmatrix} \vdots & & & & & \\ 0 & & & & & 0 \\ -t_j & \ddots & & & & \\ 0 & \ddots & \ddots & & & \\ \vdots & \ddots & \ddots & \ddots & & \\ 0 & \cdots & 0 & -t_j & 0 & \cdots \end{pmatrix} \quad \text{and } \beta_j = \begin{pmatrix} \cdots & 0 & -\frac{q_i}{i} & 0 & \cdots & 0 \\ & & \ddots & \ddots & \ddots & \vdots \\ & & & \ddots & \ddots & 0 \\ & & & & \ddots & -\frac{q_i}{i} \\ & 0 & & & & 0 \\ & & & & & \vdots \end{pmatrix},$$

$$\begin{aligned} B(\alpha_i, a_j) &= \det(I + \alpha_i^{-1} a_j^{-1} b_j \beta_i) \\ &= 1 + \sum_{n \geq 1} \text{Tr}(\Lambda^n(\alpha_i^{-1} a_j^{-1} b_j \beta_i)) \end{aligned}$$

we see that only $\text{Tr}(\alpha_i^{-1} a_j^{-1} b_j \beta_i)$ contributes to the lowest non-trivial term. For the same reason the lowest non-trivial term of $\text{Tr}(\alpha_i^{-1} a_j^{-1} b_j \beta_i)$ is equal to that of $\text{Tr}(b_j \beta_i)$. A straightforward calculation gives the required result, viz., $a_{ij} = 0$ if $i \neq j$ and $a_{ii} = 1$. We resume the results of this computation in a theorem.

Theorem 4.1 *Let ρ be the element $\exp(\sum_{j \geq 1} -\frac{q_j}{j} z^{-j})$ in $\Gamma_-(N)$. Then the tau-function corresponding to the plane ρW is given by*

$$\tau_{\rho W}(t) = \tau_{\rho \circ w \circ \alpha^{-1}}(\gamma) = \exp\left(\sum_{j \geq 1} \frac{q_j}{j} t_j\right) \tau_w(\gamma) = \exp\left(\sum_{j \geq 1} \frac{q_j}{j} t_j\right) \tau_W(t).$$

5 Geometrical Interpretation of Bäcklund–Darboux Transformations

In this section we want to interpret the trivial and elementary Bäcklund–Darboux transformations on the Segal–Wilson Grassmannian. As in section 2 we consider the flows in $\Gamma_+(N)$. First consider the trivial transformations. Let ψ_W be a wavefunction of type z^ℓ , corresponding to $W \in Gr^{(\ell)}(H)$. Then clearly multiplication by $R(z) = z^k e^{-\sum_{j=1}^{\infty} \frac{q_j}{j} z^{-j}}$ maps $\psi_W(t, z)$ into $\psi_R(t, z) = R(z) \psi_W(t, z)$. Since this has to be a Segal–Wilson wavefunction of type $z^{k+\ell}$, we have to assume that $\sum_{j=1}^{\infty} |q_j| \frac{N^j}{j} < \infty$. It is then straightforward to see that ψ_R is the wavefunction corresponding to $R(z)W \in Gr^{(k+\ell)}(H)$, hence the trivial Bäcklund–Darboux transformation maps W into $R(z)W$.

Next consider the elementary Bäcklund–Darboux transformations. The following consequence of Theorem 3.1 is more or less well known (see e.g. [10] and [2]).

Corollary 5.1 *The functions $\psi_q = q \partial q^{-1} \psi_W$ and $\psi_r = r^{-1} \partial^{-1} r \psi_W$ (resp. $\psi_q^* = -q^{-1} \partial^{-1} q \psi_W^*$ and $\psi_r^* = -r \partial r^{-1} \psi_W^*$) are (adjoint) wavefunctions of planes W_q and W_r . Moreover we have the following codimension 1 inclusions:*

$$W_q \subset W, \quad W^\perp \subset W_q^\perp, \quad W \subset W_r, \quad \text{and} \quad W_r^\perp \subset W^\perp$$

Proof. From the Sato-Wilson equations one deduces directly that for all $n \geq 1$,

$$\partial_n \psi_q = (P_q \partial^n P_q^{-1})_+ \psi_q \quad \text{and} \quad \partial_n \psi_r = (P_r \partial^n P_r^{-1})_+ \psi_r.$$

This shows the first part of the claim. Consider the following subspace in $Gr(H)$:

$$W_q = \text{the closure of } \text{Span}\{\psi_q(t, z)\}.$$

The inclusions between the spaces W and W_q follows from the first relation of (3.5) and the fact that the values of a wavefunction corresponding to an element of $Gr(H)$ are lying dense in that space. Since for a suitable γ in $\Gamma_+(N)$ the orthogonal projections of $\gamma^{-1}W_q$ on $z^\ell H_+$ resp. $\gamma^{-1}W$ on $z^{\ell+1}H_+$ have a one dimensional kernel, one obtains the codimension one result. For the inclusions between the spaces W and W_r we consider the adjoint wavefunctions $\psi_W^* = P_W^{*-1} e^{-\sum t_i z^i}$ and $\psi_r^* = -r \partial r^{-1} \psi_W^*$. Since the complex conjugate $\overline{z \psi_W^*(t, z)}$ of $z \psi_W^*(t, z)$ corresponds to the space $W^\perp$, the same argument as before shows the codimension 1 inclusion:

$$W_r^\perp := \text{the closure of } \text{Span}\{\overline{z \psi_r^*(t, z)}\} \subset W^\perp.$$

Hence $\psi_r(t, z)$ corresponds to $W_r = \text{the closure of } \text{Span}\{\psi_r(t, z)\}$. This concludes the proof of the corollary. $\square$

Now suppose we are in the situation of the above Corollary. Let $s(z) \in W \cap W_q^\perp$, then $W_q = \{w \in W \mid \langle w | s(z) \rangle = 0\}$ and

$$0 = \langle \psi_q(t, z) | s(z) \rangle = \langle \frac{\partial \psi_W(t, z)}{\partial x} - q(t)^{-1} \frac{\partial q(t)}{\partial x} \psi_W(t, z) | s(z) \rangle.$$

Hence,

$$\frac{\partial \log q}{\partial x} = \frac{\partial \log \langle \psi_W(t, z) | s(z) \rangle}{\partial x}.$$

Since both $q(t)$ and $\langle \psi_W(t, z) | s(z) \rangle$ are eigenfunctions of L that define the same Bäcklund–Darboux transformation, one deduces from Theorem 3.1 that

$$q(t) = \lambda \langle \psi_W(t, z) | s(z) \rangle \quad \text{for certain } \lambda \in \mathbb{C}^\times.$$

Now notice two things. First that we can also replace $s(z) \in W \cap W_q^\perp$ by $\tilde{s}(z) = s(z) + t(z)$ with $t(z) \in W^\perp$. And next that if $\partial + u$ is a Bäcklund–Darboux transformation which maps W into a subspace U , then for $s(z) \in W \cap U^\perp$, $q(t) = \langle \psi_W(t, z) | s(z) \rangle$ gives the eigenfunction such that $\partial + u = q \partial q^{-1}$.

For the adjoint Bäcklund–Darboux transformation, one takes $t(z) \in W^\perp \cap W_r$, then $W_r^\perp = \{w \in W^\perp \mid \langle t(z) | w \rangle = 0\}$, $\langle z \psi_W^*(t, z) | \overline{t(z)} \rangle = 0$ and in a similar way as for q one obtains that

$$r(t) = \lambda \langle z \psi_W^*(t, z) | \overline{t(z)} \rangle = \lambda \langle t(z) | \overline{z \psi_W^*(t, z)} \rangle \quad \text{for certain } \lambda \in \mathbb{C}^\times.$$

Again it is obvious that we can replace $t(z)$ by $\tilde{t}(z) = s(z) + t(z)$ with $s(z) \in W$ and that this construction gives the adjoint eigenfunction.

So we conclude that one has a geometric interpretation for the elementary Bäcklund–Darboux transformation. It has its origin on the Segal–Wilson Grassmannian, viz.,

Proposition 5.1 *Let W_q and W_r be the planes of Corollary 5.1, then*

$$q(t) = \langle \psi_W(t, z) | s(z) \rangle \quad \text{and} \quad r(t) = \langle t(z) | \overline{z\psi_W^*(t, z)} \rangle$$

for certain $s(z) \in W \cap W_q^\perp$ and $t(z) \in W^\perp \cap W_r$.

The next step in analyzing the situation in $Gr(H)$ from proposition 5.1,

$$W_q \subset W \subset W_r, \dim(W_r/W) = \dim(W/W_q) = 1,$$

is to determine the tau-function corresponding to W_q resp. W_r and their relation to that of W . Let $q(t)\partial q(t)^{-1}$ (resp. $-r(t)\partial r(t)^{-1}$) act on the wavefunction $\psi_W(t, z)$ (resp. adjoint wavefunction $\psi_W^*(t, z)$) and compare this with $\psi_q(t, z)$ (resp. $\psi_r^*(t, z)$). We thus obtain

$$\begin{aligned} zP_{W_q}(t, z) &= \frac{\partial P_W(t, z)}{\partial x} + (z - q^{-1}\frac{\partial q}{\partial x})P_W(t, z) \\ zP_{W_r}^{*-1}(t, -z) &= -\frac{\partial P_W^{*-1}(t, -z)}{\partial x} + (z + r^{-1}\frac{\partial r}{\partial x})P_W^{*-1}(t, -z) \end{aligned} \quad (5.1)$$

Comparing the coefficients of z we find that

$$\frac{\partial \log q}{\partial x} = \frac{\partial \log \tau_{W_q}/\tau_W}{\partial x} \quad \text{and} \quad \frac{\partial \log r}{\partial x} = \frac{\partial \log \tau_{W_r}/\tau_W}{\partial x}. \quad (5.2)$$

This suggests the following relations

$$\tau_{W_q}(t) = \lambda q(t)\tau_W(t) \quad \text{and} \quad \tau_{W_r}(t) = \mu r(t)\tau_W(t),$$

with $\lambda, \mu \in \mathbb{C}^\times$. Since the tau-function of a plane W in $Gr(H)$ is only determined up to a constant, one may assume that the constants λ and μ are equal to 1.

We will carry out the computations for W_r . Then that for W_q is an easy consequence of it. As above let $t(z) \in H$ be such that

$$r(t) = \langle z\psi_W^* | \overline{t(z)} \rangle = \langle t(z) | \psi_{W^\perp}(\gamma(-\bar{t}), z) \rangle.$$

If $w : z^\ell H_+ \rightarrow W$ is in $\mathfrak{P}_\ell$, then we can extend it to an embedding $w_r : z^{\ell-1} H_+ \rightarrow W_r$ by putting $w_r(z^{\ell-1}) = t(z)$ and $w_r(f) = w(f)$ for $f \in z^\ell H_+$. Then w_r belongs to $\mathfrak{P}_{\ell-1}$. Next we decompose w_r with respect to $H = z^\ell H_+ \oplus \mathbb{C}z^{\ell-1} \oplus (z^{\ell-1} H_+)^\perp$ and w with respect to $H = z^\ell H_+ \oplus (z^\ell H_+)^\perp$, then we get

$$w_r = \begin{pmatrix} w_{++} & w_{+0} \\ w_{0+} & w_{00} \\ w_{-+} & w_{-0} \end{pmatrix} \quad \text{and} \quad w = \begin{pmatrix} w_+ \\ w_- \end{pmatrix} \quad \text{with} \quad w_+ = w_{++} \quad \text{and} \quad w_- = \begin{pmatrix} w_{0+} \\ w_{-+} \end{pmatrix}.$$

We do the same with the operator γ^{-1} from $\Gamma_+(N)$. This gives you respectively

$$\gamma^{-1} = \begin{pmatrix} a_{++} & a_{+0} & a_{+-} \\ 0 & 1 & a_{0-} \\ 0 & 0 & a_{--} \end{pmatrix} = \begin{pmatrix} a_{++} & b \\ 0 & c \end{pmatrix} \quad \text{with} \quad b = (a_{+0} \ a_{+-}) \quad \text{and} \quad c = \begin{pmatrix} 1 & a_{0-} \\ 0 & a_{--} \end{pmatrix}.$$

Before computing τ_{w_r} , we notice that

$$\begin{pmatrix} a_{++} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a_{++}^{-1} & -a_{++}^{-1}a_{+0} \\ 0 & 1 \end{pmatrix} - I = \begin{pmatrix} 0 & -a_{+0} \\ 0 & 0 \end{pmatrix}$$

is a 1-dimensional operator of zero-trace. Hence we have

$$\det \left(\begin{pmatrix} a_{++} & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a_{++}^{-1} & -a_{++}^{-1}a_{+0} \\ 0 & 1 \end{pmatrix} \right) = 1,$$

so that it suffices to compute the projection of $\gamma^{-1}w_r \begin{pmatrix} a_{++} & 0 \\ 0 & 1 \end{pmatrix}^{-1}$ onto $z^{\ell-1}H_+$:

$$\begin{pmatrix} a_{++}w_+a_{++}^{-1} + bw_-a_{++}^{-1} & a_{++}w_{+0} + a_{+0}w_{00} + a_{+-}w_{-0} \\ (w_{0+} + a_{0-}w_{-+})a_{++}^{-1} & w_{00} + a_{0-}w_{-0} \end{pmatrix} = \begin{pmatrix} A & \beta \\ \alpha & \delta \end{pmatrix}.$$

As A is equal to $(\gamma^{-1}wa_{++}^{-1})_+$ and γ is chosen such that this operator is invertible, we can decompose $\begin{pmatrix} A & \beta \\ \alpha & \delta \end{pmatrix}$ as follows

$$\begin{pmatrix} A & \beta \\ \alpha & \delta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \alpha A^{-1} & 1 \end{pmatrix} \begin{pmatrix} A & 0 \\ 0 & \delta - \alpha A^{-1}\beta \end{pmatrix} \begin{pmatrix} 1 & A^{-1}\beta \\ 0 & 1 \end{pmatrix}.$$

From this we see that

$$\tau_{w_r}(\gamma) = \det \left(\begin{pmatrix} A & \beta \\ \alpha & \delta \end{pmatrix} \right) = \det(A)(\delta - \alpha A^{-1}\beta) = \tau_w(\gamma)(\delta - \alpha A^{-1}\beta).$$

Hence we merely have to show that $r(t) = \delta - \alpha A^{-1}\beta$. We know that $g(r) = \gamma(-t)^*$ with $\gamma(t) \in \Gamma_+^W(N)$ and thus we get

$$r(t) = \langle \gamma(t)t(z) | \hat{\psi}_{W^\perp}(\gamma(-t)^*, z) \rangle.$$

By definition one has

$$\hat{\psi}_{W^\perp}(\gamma(-t)^*, z) = z^{\ell-1} + \sum_{k \geq \ell} \psi_k(\gamma(-t)^*)z^k.$$

Therefore, if $\beta = \sum_{k \geq \ell} \beta_k(t)z^k$, then the expression for $r(t)$ becomes

$$\begin{aligned} r(t) &= \langle \delta z^{\ell-1} + \sum_{k \geq \ell} \beta_k(t)z^k | z^{\ell-1} + \sum_{j \geq \ell} \psi_j(\gamma(-t)^*)z^j \rangle \\ &= \delta + \sum_{k \geq \ell} \beta_k(t) \overline{\psi_k(\gamma(-t)^*)}. \end{aligned}$$

From the computation of $\hat{\psi}_{W^\perp}$ in [10], we see that, if $\alpha A^{-1}(z^{k-1}) = \alpha_k z^{\ell-1}$ for all $k \geq \ell$, then $\psi_k(\gamma(-t)^*) = -\overline{\alpha_k}$. In other words, there holds

$$r(t) = \delta - \alpha A^{-1}\beta.$$

Next we treat the case $W_q \subset W$ and $\dim(W/W_q) = 1$. Let $s(z) \in W$ be such that $W_q = \{w \in W \mid \langle w | s(z) \rangle = 0\}$ and $q(t) = \langle \psi_W(t, z) | s(z) \rangle$. Let $w_0 \in \mathfrak{P}_{\ell+1}$ be such that $w_0(z^{\ell+1}H_+) = W_q$, then $w = (w_0 | s(z))$ belongs to $\mathfrak{P}_\ell$ and $w(z^\ell H_+) = W$. According to the foregoing result, we have

$$\tau_w(t) = r(t)\tau_{w_0}(t) \quad \text{with } r(t) = \langle s(z) | \psi_{W_q^\perp}(\gamma(-t)) \rangle.$$

In particular, one has the relations

$$P_W = r^{-1}\partial^{-1}rP_{W_q} \quad \text{and} \quad P_{W_q} = q\partial q^{-1}P_W,$$

so that we may conclude

$$r^{-1}\partial r = q\partial q^{-1}.$$

According to Lemma 3.2 r^{-1} satisfies

$$\partial_n(r^{-1}) = (r^{-1}\partial^{-1}rP_{W_q}\partial^n P_{W_q}^{-1}r^{-1}\partial r)_+(r^{-1}) = (L_W^n)_+(r^{-1})$$

like q . Hence by Theorem 3.1 we may conclude that $q(t) = \lambda r(t)^{-1}$ with $\lambda \in \mathbb{C}^\times$. We resume the forgoing result in the form of a theorem.

Theorem 5.1 *Let $W, W_q, W_r \in Gr(H)$ be such that $W_q \subset W \subset W_r$ and $\dim(W_r/W) = \dim(W/W_q) = 1$. Then the tau-function of W_r is equal to $\tau_{W_r}(t) = r(t)\tau_W(t)$, with $r(t) = \langle v | \psi_{W^\perp}(\bar{t}, z) \rangle$ for some $v \in W_r \setminus W$ and the tau-function of W_q is equal to $\tau_{W_q}(t) = q(t)\tau_W(t)$, where $q(t) = \langle \psi_W(t, z) | s \rangle$ for some $s \in W \cap (W_q)^\perp$. Moreover, the wave functions are related as follows*

$$\begin{aligned} \psi_q(t, z) &= z \frac{q(t - [z^{-1}])}{q(t)} \psi_W(t, z), & \psi_q^*(t, z) &= z^{-1} \frac{q(t + [z^{-1}])}{q(t)} \psi_W^*(t, z), \\ \psi_r(t, z) &= z^{-1} \frac{r(t - [z^{-1}])}{r(t)} \psi_W(t, z), & \psi_r^*(t, z) &= z \frac{r(t + [z^{-1}])}{r(t)} \psi_W^*(t, z). \end{aligned} \tag{5.3}$$

6 Squared Eigenfunction Potentials

Let q be an eigenfunction and r be an adjoint eigenfunction of the KP Lax operator L . For such a pair of (adjoint) eigenfunctions Oevel [17] showed that there exists a function $\Omega(r, q)$ called the *squared eigenfunction potential*, which satisfies

$$\frac{\partial \Omega(r, q)}{\partial t_k} = \text{Res}_\partial(\partial^{-1}r(L^k)_+ q \partial^{-1}). \tag{6.1}$$

This potential was described in [15] and [7] (see also [17] and [18]). A special case of (6.1) is the case that $k = 1$, i.e. $t_1 = x$, then

$$\frac{\partial \Omega(r, q)}{\partial x} = qr.$$

Equation (6.1) determines Ω uniquely upto a shift by a constant.

The expressions $\partial^{-1}r\psi$ and $\partial^{-1}\psi^*q$ are special squared eigenfunction potentials. Notice that they are uniquely defined. We now want to find an explicit expression for these specific potentials in the Segal-Wilson setting. Since

$$\psi_r = r^{-1}\partial^{-1}r\psi_W \quad \text{and} \quad \psi_q^* = -q^{-1}\partial^{-1}q\psi_W,$$

one has

$$\begin{aligned} \partial^{-1}r(t)\psi_W(t, z) &= r(t)\psi_r(t, z) \\ &= r(t)z^{-1} \frac{r(t - [z^{-1}])}{r(t)} \psi_W(t, z) \\ &= z^{-1}r(t - [z^{-1}])\psi_W(t, z) \end{aligned}$$

and

$$\begin{aligned}\partial^{-1}q(t)\psi_W^*(t, z) &= -q(t)\psi_q^*(t, z) \\ &= -q(t)z^{-1}\frac{q(t + [z^{-1}])}{q(t)}\psi_W^*(t, z) \\ &= -z^{-1}q(t + [z^{-1}])\psi_W^*(t, z).\end{aligned}$$

We can also express them in $s(z)$. Let

$$q(t) = \langle \psi_W(t, z) | s(z) \rangle, \quad \text{and} \quad r(t) = \langle \psi_W^*(t, z) | \overline{zs(z)} \rangle,$$

then $\partial^{-1}r\psi$ and $\partial^{-1}\psi^*q$ are equal to

$$\begin{aligned}S(\psi_W(t, z), \overline{zs(z)}) &:= z^{-1} \langle \psi_W^*(t - [z^{-1}], y) | \overline{ys(y)} \rangle \psi_W(t, z), \\ S(\psi_W^*(t, z), s(z)) &:= -z^{-1} \psi_W^*(t, z) \langle \psi_W(t + [z^{-1}], y) | s(y) \rangle,\end{aligned}\tag{6.2}$$

respectively.

Next let $W \in Gr^{(\ell)}(H)$, then, since $\psi_W(t, z)$ and $\psi_W^*(t, z)$ satisfy (1.13) we find the following expressions for these squared eigenfunction potentials (see also [4]).

Proposition 6.1 *Let $|z| > |y|$, then for $W \in Gr^{(\ell)}(H)$ on has*

$$\begin{aligned}S(\psi_W(t, z), \overline{zs(z)}) &= \frac{\langle (z/y)^\ell X(t, z, y) \tau_W(t) | \overline{ys(y)} \rangle}{\tau_W(t)}, \\ S(\psi_W^*(t, z), s(z)) &= \frac{\langle (y/z)^\ell X(t, y, z) \tau_W(t) | s(y) \rangle}{\tau_W(t)},\end{aligned}\tag{6.3}$$

where

$$X(t, \lambda, \mu) = \frac{1}{\lambda - \mu} \exp\left(\sum t_i(\lambda^i - \mu^i)\right) \exp\left(\sum \frac{\mu^{-i} - \lambda^{-i}}{i} \frac{\partial}{\partial t_i}\right)\tag{6.4}$$

is the vertex operator for the KP hierarchy.

This vertex operator expresses the action of the Lie algebra gl_∞ , see [14] for more details.

The results of this section, will be used in the following section on geometric Bäcklund–Darboux transformations.

7 Geometric Bäcklund–Darboux Transformations

We can of course take the converse of the procedure of section 5 and use the geometry to construct elementary Bäcklund–Darboux transformations, viz as follows:

Theorem 7.1 *Let $W \in Gr(H)$ with τ_W its tau-function, ψ_W its wavefunction and ψ_W^* its adjoint wavefunction. Let $s(z) \in H$, then $q(t) = \langle \psi_W(t, z) | s(z) \rangle$ is an eigenfunction and $r(t) = \langle s(z) | \psi_W^*(t, z) \rangle$ is an adjoint eigenfunction and the corresponding elementary Bäcklund–Darboux transformations $q\partial q^{-1}$ and $r^{-1}\partial^{-1}r$ give new tau- and wavefunctions $\tau_{W_q}(t) = q(t)\tau_W(t)$, $\tau_{W_r}(t) = r(t)\tau_W(t)$, $\psi_q = q\partial q^{-1}\psi_W$, $\psi_r = r^{-1}\partial^{-1}r\psi_W$ and adjoint wave functions $\psi_q^* = -q^{-1}\partial^{-1}q\psi_W^*$, $\psi_r^* = -r\partial r^{-1}\psi_W^*$ corresponding to planes $W_q = \{w \in W | \langle w | s(z) \rangle = 0\}$, $W_r = W \oplus \mathbb{C}s(z)$, $W_q^\perp = W^\perp \oplus \mathbb{C}s(z)$, $W_r^\perp = \{w \in W^\perp | \langle s(z) | w \rangle = 0\}$, respectively.*

We now want to generalize the situation of the previous Theorem. Assume that is given the plane $W \in Gr^{(\ell)}(H)$ with wavefunction $\psi_W(t, z)$ and adjoint wavefunction $\psi_W^*(t, z)$. We want to determine the Bäcklund–Darboux transformation $BD(V, W)$ that maps $\psi_W(t, z)$ to $\psi_V(t, z)$. We restrict ourselves to the case that $BD(V, W)$ is a finite product of elementary Bäcklund–Darboux transformation, i.e., to the case that $V \cap W$ has finite codimension inside both V and W . Of course one has that if $\psi_W(t, z) = P_W(\partial)e^{\sum t_i z^i}$ and $\psi_V(t, z) = P_V(\partial)e^{\sum t_i z^i}$, that $BD(V, W) = P_V(\partial)P_W(\partial)^{-1}$. But we assume that the wavefunction and its adjoint are only given for W and not yet known for V .

We first start with two special cases of planes V and W , viz., $V \subset W$ and $W \subset V$. First consider the former case and assume that V has codimension n inside W . So we can find n independent functions in $W \cap V^\perp$, say $w_1, w_2, \dots, w_n$. It is straightforward that $BD(V, W)$ is the product of n elementary Bäcklund–Darboux transformations of the form $q_j \partial q_j^{-1}$. Hence, $BD(V, W) = \sum_{k=0}^n a_k \partial^k$ with $a_k \in R$ and $a_n = 1$, i.e. is a differential operator of order n with leading coefficient equal to 1. Thus, $\psi_V = \sum_{k=0}^n a_k \frac{\partial^k \psi_W}{\partial x^k}$ and since $\langle \psi_V | w_j \rangle = 0$ for all $j = 1, 2, \dots, n$, we find that

$$\sum_{k=0}^{n-1} a_k \langle \frac{\partial^k \psi_W(t, z)}{\partial x^k} | w_j(z) \rangle = - \langle \frac{\partial^n \psi_W(t, z)}{\partial x^n} | w_j(z) \rangle, \quad \text{for } j = 1, 2, \dots, n.$$

Using Cramer's rule one thus finds the following expressions for the a_k 's:

$$a_k = (-)^{n-k} \mathcal{W}(\psi_W; w_1, w_2, \dots, w_n)^{-1} \mathcal{W}_k(\psi_W; w_1, w_2, \dots, w_n),$$

where

$$\mathcal{W}(A; w_1, w_2, \dots, w_n) = \det \begin{pmatrix} \langle A | w_1 \rangle & \langle \frac{\partial A}{\partial x} | w_1 \rangle & \cdots & \langle \frac{\partial^{n-1} A}{\partial x^{n-1}} | w_1 \rangle \\ \langle A | w_2 \rangle & \langle \frac{\partial A}{\partial x} | w_2 \rangle & \cdots & \langle \frac{\partial^{n-1} A}{\partial x^{n-1}} | w_2 \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle A | w_n \rangle & \langle \frac{\partial A}{\partial x} | w_n \rangle & \cdots & \langle \frac{\partial^{n-1} A}{\partial x^{n-1}} | w_n \rangle \end{pmatrix} \quad (7.1)$$

is a Wronskian determinant and

$$\mathcal{W}_k(A; w_1, w_2, \dots, w_n) = \det \begin{pmatrix} \langle A | w_1 \rangle & \cdots & \langle \frac{\partial^{k-1} A}{\partial x^{k-1}} | w_1 \rangle & \langle \frac{\partial^{k+1} A}{\partial x^{k+1}} | w_1 \rangle & \cdots & \langle \frac{\partial^n A}{\partial x^n} | w_1 \rangle \\ \langle A | w_2 \rangle & \cdots & \langle \frac{\partial^{k-1} A}{\partial x^{k-1}} | w_2 \rangle & \langle \frac{\partial^{k+1} A}{\partial x^{k+1}} | w_2 \rangle & \cdots & \langle \frac{\partial^n A}{\partial x^n} | w_2 \rangle \\ \vdots & & \vdots & \vdots & & \vdots \\ \langle A | w_n \rangle & \cdots & \langle \frac{\partial^{k-1} A}{\partial x^{k-1}} | w_n \rangle & \langle \frac{\partial^{k+1} A}{\partial x^{k+1}} | w_n \rangle & \cdots & \langle \frac{\partial^n A}{\partial x^n} | w_n \rangle \end{pmatrix}.$$

This is in fact a generalization of a result of Crumm [8]. Following [17], we give a short notation for this Bäcklund–Darboux transformation $BD(V, W)$, viz.,

$$BD(V, W) = \mathcal{W}(\psi_W; w_1, w_2, \dots, w_n)^{-1} \mathcal{W}(\psi_W; w_1, w_2, \dots, w_n; \partial), \quad (7.2)$$

where

$$\mathcal{W}(A; w_1, w_2, \dots, w_n; \partial) = \det \begin{pmatrix} \langle A|w_1 \rangle & \langle \frac{\partial A}{\partial x}|w_1 \rangle & \cdots & \langle \frac{\partial^n A}{\partial x^n}|w_1 \rangle \\ \langle A|w_2 \rangle & \langle \frac{\partial A}{\partial x}|w_2 \rangle & \cdots & \langle \frac{\partial^n A}{\partial x^n}|w_2 \rangle \\ \vdots & \vdots & \ddots & \vdots \\ \langle A|w_n \rangle & \langle \frac{\partial A}{\partial x}|w_n \rangle & \cdots & \langle \frac{\partial^n A}{\partial x^n}|w_n \rangle \\ 1 & \partial & \cdots & \partial^n \end{pmatrix}. \quad (7.3)$$

The determinant (7.3) must be seen as a formal expansion with respect to the last row vector, collecting all minors left of the differential symbols ∂^k .

It is obvious that the Bäcklund-Darboux transformation $BD(V, W)$ does not depend on the choice of basis of the linear space $W \cap V^\perp$. It is also obvious that one can even add elements of $W^\perp$ to the w_k 's.

We now want to determine the inverse of this Bäcklund-Darboux transformation $BD(V, W)$. We immitate the proof of the Lemma in [19]. Let b_k , $1 \leq k \leq n$ be the solution of the linear system

$$\mathcal{W}(\psi_W; w_1, w_2, \dots, w_n) b_k = \delta_{kn},$$

i.e.,

$$b_k = (-)^{n-k} \mathcal{W}(\psi_W; w_1, w_2, \dots, w_n)^{-1} \mathcal{W}(\psi_W; w_1, w_2, \dots, w_{k-1}, w_{k+1}, \dots, w_n)$$

then

$$BD(V, W)^{-1} = \sum_{k=1}^n \langle \psi_W | w_k \rangle \partial^{-1} b_k.$$

To show this we use the fact that for $f \in R$

$$f \partial^{-1} = -(\partial^{-1} f)^* = \sum_{\ell=0}^{\infty} \left((-)^{\ell+1} \frac{\partial^\ell f}{\partial x^\ell} \partial^{-\ell-1} \right)^* = \sum_{\ell=0}^{\infty} \partial^{-\ell-1} \frac{\partial^\ell f}{\partial x^\ell}$$

and calculate

$$\begin{aligned} BD(V, W) \sum_{k=1}^n \langle \psi_W | w_k \rangle \partial^{-1} b_k &= (BD(V, W) \sum_{k=1}^n \langle \psi_W | w_k \rangle \partial^{-1} b_k)_+ + (BD(V, W) \sum_{k=1}^n \langle \psi_W | w_k \rangle \partial^{-1} b_k)_- \\ &= (BD(V, W) \sum_{\ell=0}^{\infty} \partial^{-\ell-1} \sum_{k=1}^n \langle \frac{\partial^\ell \psi_W}{\partial x^\ell} | w_k \rangle \partial^{-1} b_k)_+ + \sum_{k=1}^n BD(V, W) (\langle \psi_W | w_k \rangle) \partial^{-1} b_k \\ &= (BD(V, W) \sum_{\ell=0}^{n-1} \partial^{-\ell-1} \mathcal{W}(\psi_W; w_1, w_2, \dots, w_n))_+ + 0 \\ &= (BD(V, W) \partial^{-n})_+ \\ &= 1. \end{aligned}$$

Again one has a short notation for

$$BD(V, W)^{-1} = \mathcal{W}(\partial; \psi_W; w_1, w_2, \dots, w_n) \mathcal{W}(\psi_W; w_1, w_2, \dots, w_n)^{-1}, \quad (7.4)$$

where

$$\mathcal{W}(\partial; A; w_1, w_2, \dots, w_n) = \det \begin{pmatrix} \langle A|w_1 \rangle & \langle \frac{\partial A}{\partial x}|w_1 \rangle & \cdots & \langle \frac{\partial^{n-1} A}{\partial x^{n-2}}|w_1 \rangle & \langle A|w_1 \rangle \partial^{-1} \\ \langle A|w_2 \rangle & \langle \frac{\partial A}{\partial x}|w_2 \rangle & \cdots & \langle \frac{\partial^{n-1} A}{\partial x^{n-2}}|w_2 \rangle & \langle A|w_2 \rangle \partial^{-1} \\ \vdots & \vdots & \cdots & \vdots & \vdots \\ \langle A|w_n \rangle & \langle \frac{\partial A}{\partial x}|w_n \rangle & \cdots & \langle \frac{\partial^{n-1} A}{\partial x^{n-2}}|w_n \rangle & \langle A|w_n \rangle \partial^{-1} \end{pmatrix} \quad (7.5)$$

In (7.5) one has to use the expansion of the determinant with respect to the last column collecting all minors on the right of the $\langle A|w_k \rangle \partial^{-1}$ symbols.

Next consider the other special case, viz., that $W \subset V$ of codimension m . Then clearly $BD(V, W)$ is the product of m elementary adjoint Bäcklund-Darboux transformations $r_i^{-1} \partial^{-1} r_i$. We first do not consider $BD(V, W)$ but rather $BD(V, W)^{* -1}$, which is a differential operator of order m , say $\sum_{k=0}^m a_k \partial^k$. Then notice that $a_m = (-1)^m$ and $\psi_V^* = \sum_{k=0}^m a_k \frac{\partial^k \psi_W^*}{\partial x^k}$. Now let $v_1, v_2, \dots, v_m$ be m linearly independent elements in $W^\perp \cap V$. For notational convenience we introduce $v^\dagger$ by

$$v^\dagger(z) := \overline{zv(z)}. \quad (7.6)$$

Then $\langle \psi_V^* | v_j^\dagger \rangle = 0$ and hence

$$\sum_{k=0}^{m-1} a_k \langle \frac{\partial^k \psi_W^*}{\partial x^k} | v_j^\dagger \rangle = (-1)^{m+1} \langle \frac{\partial^m \psi_W^*}{\partial x^m} | v_j^\dagger \rangle, \quad \text{for } j = 1, 2, \dots, m.$$

Using Cramer's rule again one obtains the following expression for the a_k 's:

$$a_k = (-1)^k \mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger)^{-1} \mathcal{W}_k(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger).$$

In a similar way as before one finds

$$BD(V, W)^{* -1} = (-1)^m \mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger)^{-1} \mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger; \partial), \quad (7.7)$$

where one has to use the expansion of $\mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger; \partial)$ with respect to the last row collecting all minors on the left of the ∂ symbols. So

$$BD(V, W)^{-1} = (-1)^m \mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger; -\partial) \mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger)^{-1} \quad (7.8)$$

and here one has to use the expansion of $\mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger; -\partial)$ with respect to the last row collecting all minors on the right of the ∂ symbols.

In a similar way as before we now determine $BD(V, W)^*$. Let b_k , $1 \leq k \leq m$ be a solution of the linear system

$$\mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger) b_k = (-1)^m \delta_{mk},$$

then

$$BD(V, W)^* = \sum_{k=1}^m \langle \psi_W^* | v_k^\dagger \rangle \partial^{-1} b_k,$$

with

$$b_k = (-)^k \mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger)^{-1} \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_{k-1}^\dagger, v_{k+1}^\dagger, \dots, v_m^\dagger).$$

So

$$BD(V, W) = - \sum_{k=1}^m b_k \partial^{-1} \langle \psi_W^* | v_k^\dagger \rangle$$

and hence

$$BD(V, W) = (-)^m \mathcal{W}(\psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger)^{-1} \mathcal{W}(\partial; \psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger)^*, \quad (7.9)$$

where one has to use the expansion of the determinant of $\mathcal{W}(\partial; \psi_W^*; v_1^\dagger, v_2^\dagger, \dots, v_m^\dagger)$ with respect to the last column collecting all minors on the right of the $\langle \psi_W^* | v_k^\dagger \rangle \partial^{-1}$ symbols. Again also in this case it is clear that everything is independent from the choice of basis of $W^\perp \cap V$.

We now use the results of the previous two cases to determine the Bäcklund-Darboux transformation $BD(V, W)$ in the more general case that $V \cap W$ has codimension m in V and codimension n in W . Let $w_1, w_2, \dots, w_n$ be a basis of $V^\perp \cap W$ and let $v_1, v_2, \dots, v_m$ be a basis of $W^\perp \cap V$. Starting with W we first apply m elementary adjoint Bäcklund-Darboux transformations to obtain $W+V$ and then apply n elementary Bäcklund-Darboux transformations to obtain V . In other words

$$BD(V, W) = BD(V, V+W)BD(V+W, W).$$

Now $BD(V, V+W)$ is a differential operator of order n for which the leading coefficient is equal to 1 and $BD(V+W, W) = \sum_{k=1}^m b_k \partial^{-1} \langle \psi_W^* | v_k^\dagger \rangle$ and is in fact a pseudodifferential operator of order $-m$, again with leading coefficient equal to 1. Hence

$$\begin{aligned} BD(V, W) &= \left(\sum_{\ell=0}^n c_\ell \partial^\ell \right) \left(\sum_{k=1}^m b_k \partial^{-1} \langle \psi_W^* | v_k^\dagger \rangle \right) \\ &= \sum_{k=1}^m a_{-k} \partial^{-1} \langle \psi_W^* | v_k^\dagger \rangle + \sum_{\ell=0}^{n-m} a_\ell \partial^\ell \end{aligned}$$

is a pseudodifferential operator of order $n-m$ with leading coefficient equal to 1 (notice that this is a_{n-m} if $n \geq m$). Since $BD(V, W)\psi_W$ belongs to V it must be perpendicular to $w_1, w_2, \dots, w_n$, which gives the restriction

$$\sum_{k=1}^m a_{-k} S(\psi_W, v_k^\dagger, w_j) + \sum_{\ell=0}^{n-m} a_\ell \langle \frac{\partial^\ell \psi_W}{\partial x^\ell} | w_j \rangle = 0, \quad (7.10)$$

Where (cf. (6.2))

$$S(A, s, t) := \langle S(A, s) | t \rangle. \quad (7.11)$$

Now if $n \geq m$, the equations (7.10) determine $BD(V, W)$, since $a_{n-m} = 1$. However, if $n < m$ then (7.10) is equal to

$$\sum_{k=1}^m a_{-k} S(\psi_W, v_k^\dagger, w_j) = 0, \quad (7.12)$$

and we need some more information. We also know that $BD(V, W)$ is a pseudodifferential operator of order $n - m$ with leading coefficient equal to 1. This leads to the extra restrictions.

$$\sum_{k=1}^m a_{-k} \left\langle \frac{\partial^j \psi_W^*}{\partial x^j} | v_k^\dagger \right\rangle = (-)^{m-n-1} \delta_{j, m-n-1} \quad \text{for } j = 0, 1, \dots, m-n-1. \quad (7.13)$$

Then (7.12) and (7.13) determine $BD(V, W)$ uniquely. Using Cramer's rule we can determine in the case that $n \geq m$ from (7.10) and in the case that $m > n$ from (7.12) and (7.13) the coefficients a_i of $BD(V, W)$. We thus obtain

$$BD(V, W) = \mathcal{W}(\psi_W; w_1, \dots, w_n; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger)^{-1} \mathcal{W}(\psi_W; w_1, \dots, w_n; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \partial), \quad (7.14)$$

where

$$\mathcal{W}(A; w_1, \dots, w_n; B; v_1, \dots, v_m) = \det \begin{pmatrix} S(A, v_1, w_1) & \cdots & S(A, v_m, w_1) & \langle A | w_1 \rangle & \cdots & \langle \frac{\partial^{n-m-1} A}{\partial x^{n-m-1}} | w_1 \rangle \\ \vdots & & \vdots & \vdots & & \vdots \\ S(A, v_1, w_n) & \cdots & S(A, v_m, w_n) & \langle A | w_n \rangle & \cdots & \langle \frac{\partial^{n-m-1} A}{\partial x^{n-m-1}} | w_n \rangle \\ \langle B | v_1 \rangle & \cdots & \langle B | v_m \rangle & & & \\ \langle \frac{\partial B}{\partial x} | v_1 \rangle & \cdots & \langle \frac{\partial B}{\partial x} | v_m \rangle & & & \\ \vdots & & \vdots & & & \\ \langle \frac{\partial^{m-n-1} B}{\partial x^{m-n-1}} | v_1 \rangle & \cdots & \langle \frac{\partial^{m-n-1} B}{\partial x^{m-n-1}} | v_m \rangle & & & \end{pmatrix} \quad (7.15)$$

and

$$\mathcal{W}(A; w_1, \dots, w_n; B; v_1, \dots, v_m; \partial) = \det \begin{pmatrix} S(A, v_1, w_1) & \cdots & S(A, v_m, w_1) & \langle A | w_1 \rangle & \cdots & \langle \frac{\partial^{n-m} A}{\partial x^{n-m}} | w_1 \rangle \\ \vdots & & \vdots & \vdots & & \vdots \\ S(A, v_1, w_n) & \cdots & S(A, v_m, w_n) & \langle A | w_n \rangle & \cdots & \langle \frac{\partial^{n-m} A}{\partial x^{n-m}} | w_n \rangle \\ \partial^{-1} \langle B | v_1 \rangle & \cdots & \partial^{-1} \langle B | v_m \rangle & 1 & \cdots & \partial^{n-m} \\ \langle B | v_1 \rangle & \cdots & \langle B | v_m \rangle & & & \\ \langle \frac{\partial B}{\partial x} | v_1 \rangle & \cdots & \langle \frac{\partial B}{\partial x} | v_m \rangle & & & \\ \vdots & & \vdots & & & \\ \langle \frac{\partial^{m-n-2} B}{\partial x^{m-n-2}} | v_1 \rangle & \cdots & \langle \frac{\partial^{m-n-2} B}{\partial x^{m-n-2}} | v_m \rangle & & & \end{pmatrix}. \quad (7.16)$$

The matrix appearing in (7.15) is an $n \times n$ -matrix if $n \geq m$ and an $m \times m$ -matrix if $n < m$. The matrix appearing in (7.16) is an $n+1 \times n+1$ -matrix if $n \geq m$ and an $m \times m$ -matrix if $n < m$. The determinant appearing in (7.16) is the formal expansion with respect to the $(n+1)$ -th row putting all the minors to the left of the ∂ symbols.

Finally we want to calculate the inverse of $BD(V, W)$ or rather the adjoint of the inverse. First notice that

$$\begin{aligned} BD(V, W)^{* -1} &= BD(V^\perp, W^\perp) \quad \text{and} \\ BD(V, W)^{* -1} &= BD(V, V \cap W^{* -1} BD(V \cap W, W)^{* -1}). \end{aligned} \quad (7.17)$$

From the first line of (7.17) we deduce that

$$\langle BD(V, W)^{* -1} \psi_W^* | v_j^\dagger \rangle = 0 \quad \text{for } j = 1, 2, \dots, m$$

and from the second line that

$$BD(V, W)^{* -1} = \sum_{\ell=1}^n a_{-\ell} \partial^{-1} \langle \psi_W | w_\ell \rangle + \sum_{k=0}^{m-n} a_k \partial^k.$$

Using the fact that $BD(V, W)^{* -1}$ is an $(m-n)$ -th order pseudodifferential operator with leading coefficient $(-1)^{m-n}$ we obtain that for $1 \leq j \leq m$

$$\sum_{\ell=1}^n a_{-\ell} S(\psi_W^*, w_\ell, v_j^\dagger) + \sum_{k=0}^{m-n-1} a_k \langle \frac{\partial^k \psi_W^*}{\partial x^k} | v_j^\dagger \rangle = (-1)^{n-m-1}, \quad (7.18)$$

if $m \geq n$ and that if $m < n$

$$\begin{aligned} \sum_{\ell=1}^n a_{-\ell} S(\psi_W^*, w_\ell, v_j^\dagger) &= 0, \\ \sum_{\ell=1}^n a_{-\ell} \langle \frac{\partial^k \psi_W}{\partial x^k} | w_\ell \rangle &= -\delta_{k, n-m-1}, \end{aligned} \quad (7.19)$$

for $1 \leq j \leq m$ and $0 \leq k \leq n-m-1$. This determines $BD(V, W)^{* -1}$ uniquely. One has

$$\begin{aligned} BD(V, W)^{* -1} &= (-1)^{n+m} \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W; w_1, \dots, w_n)^{-1} \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W; w_1, \dots, w_n; \partial), \end{aligned} \quad (7.20)$$

hence the adjoint of this operator is

$$\begin{aligned} BD(V, W)^{-1} &= \\ (-1)^{n+m} \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W; w_1, \dots, w_n; \partial)^* \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W; w_1, \dots, w_n)^{-1}, \end{aligned} \quad (7.21)$$

We now want to calculate τ_V . In order to do that we write $BD(V, W)$ as product of $n+m$ elementary Bäcklund-Darboux transformations

$$BD(V, W) = BD(V, U_{n+m-1}) BD(U_{n+m-1}, U_{n+m-2}) \cdots BD(U_1, W),$$

where

$$U_k = \begin{cases} W + \sum_{j=1}^k \mathbb{C}v_j & \text{for } 1 \leq k \leq m, \\ \{u \in W + V \mid \langle u | w_j \rangle = 0 \text{ for } 1 \leq j \leq k - n\} & \text{for } m \leq k \leq m + n. \end{cases}$$

Then from Theorem 7.1 we deduce that for $k \leq m$

$$\begin{aligned} \tau_{U_k} &= \langle \psi_{U_{k-1}}^* | v_k^\dagger \rangle \tau_{U_{k-1}} \\ &= \langle BD(U_{k-1}, W)^{* -1} \psi_W^* | v_k^\dagger \rangle \tau_{U_{k-1}} \\ &= (-)^{k-1} \frac{\mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_k^\dagger)}{\mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_{k-1}^\dagger)} \tau_{U_{k-1}} \end{aligned}$$

and that for $k > m$

$$\begin{aligned} \tau_{U_k} &= \langle \psi_{U_{k-1}} | w_{k-m} \rangle \tau_{U_{k-1}} \\ &= \langle BD(U_{k-1}, W) \psi_W | w_{k-m} \rangle \tau_{U_{k-1}} \\ &= \frac{\mathcal{W}(\psi_W; w_1, \dots, w_{k-m}; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger)}{\mathcal{W}(\psi_W; w_1, \dots, w_{k-m-1}; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger)} \tau_{U_{k-1}}. \end{aligned}$$

From which we deduce that (upto a scalar multiple)

$$\tau_V = \mathcal{W}(\psi_W; w_1, \dots, w_n; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger) \tau_W. \quad (7.22)$$

Notice that all the above considerations are independent of the choice of bases of $V \cap W^\perp$ and $V^\perp \cap W$. Thus we have proven the main theorem of this paper

Theorem 7.2 *Let $V \in Gr^{(k)}$ and $W \in Gr^{(\ell)}$ be such that $V \cap W$ has codimension m in V and codimension n in W (hence $k - m = \ell - n$) and let ψ_W , ψ_W^* and τ_W be the to W corresponding wave-, adjoint wave- and tau-function. Then the corresponding Bäcklund-Darboux transformation is equal to (cf. (7.15), (7.16))*

$$BD(V, W) = \mathcal{W}(\psi_W; w_1, \dots, w_n; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger)^{-1} \mathcal{W}(\psi_W; w_1, \dots, w_n; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \partial),$$

and its inverse is equal to

$$\begin{aligned} BD(V, W)^{-1} &= \\ &(-)^{n+m} \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W; w_1, \dots, w_n; \partial)^* \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W; w_1, \dots, w_n)^{-1} \end{aligned}$$

for any bases $w_1, w_2, \dots, w_n$ of $W \cap V^\perp$ and $v_1, v_2, \dots, v_m$ of $V \cap W^\perp$. Moreover,

$$\begin{aligned} \psi_V &= \mathcal{W}(\psi_W; w_1, \dots, w_n; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger)^{-1} \mathcal{W}(\psi_W; w_1, \dots, w_n; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W), \\ \psi_V^* &= \\ &(-)^{n+m} \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W; w_1, \dots, w_n)^{-1} \mathcal{W}(\psi_W^*; v_1^\dagger, \dots, v_m^\dagger; \psi_W; w_1, \dots, w_n; \psi_W^*), \end{aligned}$$

and

$$\tau_V = \mathcal{W}(\psi_W; w_1, \dots, w_n; \psi_W^*; v_1^\dagger, \dots, v_m^\dagger) \tau_W,$$

where

$$\mathcal{W}(A; w_1, \dots, w_n; B; v_1, \dots, v_m; C) = \det \begin{pmatrix} S(A, v_1, w_1) & \cdots & S(A, v_m, w_1) & \langle A|w_1 \rangle & \cdots & \langle \frac{\partial^{n-m} A}{\partial x^{n-m}} | w_1 \rangle \\ \vdots & & \vdots & \vdots & & \vdots \\ S(A, v_1, w_n) & \cdots & S(A, v_m, w_n) & \langle A|w_n \rangle & \cdots & \langle \frac{\partial^{n-m} A}{\partial x^{n-m}} | w_n \rangle \\ S(C, v_1) & \cdots & S(C, v_m) & C & \cdots & \frac{\partial^{n-m} C}{\partial x^{n-m}} \\ \langle B|v_1 \rangle & \cdots & \langle B|v_m \rangle & & & \\ & \cdots & \langle \frac{\partial B}{\partial x} | v_m \rangle & & & \\ \vdots & & \vdots & & & \\ \langle \frac{\partial^{m-n-2} B}{\partial x^{m-n-2}} | v_1 \rangle & \cdots & \langle \frac{\partial^{m-n-2} B}{\partial x^{m-n-2}} | v_m \rangle & & & \end{pmatrix}, \quad (7.23)$$

where $S(C, v) = \partial^{-1}(\langle B|v \rangle C)$ is defined in (6.2) and $S(A, v, w)$ in (7.11).

8 Bäcklund–Darboux Transformations for the Gelfand–Dickey Hierarchy

In this section we want to consider the elementary Bäcklund–Darboux transformations for the Gelfand–Dickey hierarchies. It is well-known that these correspond to the n -th reduced KP hierarchy, i.e., $L^n = (L^n)_+$. Since $L^n \psi_W(t, z) = z^n \psi_W(t, z)$ we have a trivial Bäcklund–Darboux transformation that maps W into $z^n W$. Since $L^n = (L^n)_+$, $z^n \psi_W(t, z) = \partial_n(\psi_W(t, z))$ and hence $z^n W \subset W$. We now want to determine which vectors $s \in W$ we can choose such that $W_s = \{w \in W \mid \langle w|s \rangle = 0\}$ also satisfies $z^n W_s \subset W_s$.

We will first prove the following Lemma.

Lemma 8.1 *Let $W' \subset W$ such that $z^n W' \subset W$ and*

$$W_s = \{w \in W \mid \langle w|s \rangle = 0\} \quad \text{and} \quad W'_s = \{w \in W' \mid \langle w|s \rangle = 0\}.$$

Then $z^n W'_s \subset W_s$ if and only if there exists a $\lambda \in \mathbb{C}$ such that

$$(z^{-n} - \lambda)s \in W'^{\perp}. \quad (8.1)$$

Proof. Let $v \in W'_s$ and $z^n v \in W_s$, then

$$\langle z^n v|s(z) \rangle = \langle v|z^{-n} s(z) \rangle = 0,$$

in other words $z^{-n} s(z) \in W_s'^{\perp}$. Hence, since $W_s'^{\perp} = W'^{\perp} \oplus \mathbb{C}s$, there must exist a $\lambda \in \mathbb{C}$ such that $(z^{-n} - \lambda)s \in W'^{\perp}$.

Now assume that (8.1) holds, then for every $v \in W'_s$ one has

$$\langle z^n v|s(z) \rangle = \langle v|z^{-n} s(z) \rangle = \bar{\lambda} \langle v|s(z) \rangle = 0,$$

hence $z^n v \in W_s$. □

So it suffices to assume that $W = W'$ and that $s(z) \in W$ satisfies (8.1). Let

$$U = W \cap (z^n W)^\perp, \quad (8.2)$$

then (see e.g. [21]) $W = \bigoplus_{k=0}^{\infty} z^{kn} U$, so we write

$$s(z) = \sum_{k=0}^{\infty} u_k z^{kn} \quad \text{with } u_k \in U.$$

Since $(z^{-n} - \lambda)s(z) \in W^\perp$, one obtains that $u_{k+1} = \lambda u_k$ for $k = 0, 1, \dots$ and hence,

$$s(z) = \sum_{k=0}^{\infty} (\lambda z)^{kn} u_0 \quad \text{with } u_0 \in U \text{ and } \lambda \in \mathbb{C}.$$

Since

$$\begin{aligned} \langle s(z) | s(z) \rangle &= \langle \sum_{k=0}^{\infty} (\lambda z)^{kn} u_0 | \sum_{k=0}^{\infty} (\lambda z)^{kn} u_0 \rangle \\ &= \sum_{k=0}^{\infty} |\lambda|^{2k} |u_0|^2, \end{aligned}$$

must be finite, we must choose $|\lambda| < 1$. Notice that if W satisfies $z^n W \subset W$, then $W^\perp$ satisfies $z^{-n} W^\perp \subset W^\perp$. Now let $V = W^\perp \cap (z^{-n} W^\perp)^\perp$, then $V = z^{-n} U$ and hence one obtains that $t(z) \in V$ which gives the correct Bäcklund–Darboux transformation is

$$t(z) = \sum_{k=0}^{\infty} \left(\frac{\lambda}{z}\right)^{nk} z^{-n} u_0, \quad \text{with } u_0 \in U \text{ and } \lambda \in \mathbb{C}, |\lambda| < 1.$$

We thus have proven the following theorem

Theorem 8.1 *Let $W \in Gr(H)$ satisfy $z^n W \subset W$ and hence $z^{-n} W^\perp \subset W^\perp$, then $s(z) \in W$ (resp. $t(z) \in W^\perp$) defines an eigenfunction $q(t) = \langle \psi_w(t, z) | s(z) \rangle$ (resp. adjointed eigenfunction $r(t) = \langle s(z) | z \psi_W^*(t, z) \rangle$) and $\psi_q(t, z) = q \partial q^{-1} \psi_W(t, z)$ (resp. $\psi_r(t, z) = r^{-1} \partial^{-1} r \psi_W(t, z)$) be a new wavefunction defined by the elementary Bäcklund–Darboux transformation that maps W into $W_q = \{w \in W | \langle w | s(z) \rangle = 0\}$ (resp. $W_r = W \oplus \mathbb{C} t(z)$). Let $U = W \cap (z^n W)^\perp$. Then W_q satisfies $z^n W_q \subset W_q$ (resp. W_r satisfies $z^n W_r \subset W_r$) if and only if*

$$s(z) = \sum_{k=0}^{\infty} (\lambda z)^{kn} u_0 \quad (\text{resp. } t(z) = \sum_{k=0}^{\infty} \left(\frac{\lambda}{z}\right)^{nk} z^{-n} u_0),$$

with $u_0 \in U$ and $\lambda \in \mathbb{C}$, $|\lambda| < 1$.

Next we want to give the geometrical interpretation (in terms of the Segal–Wilson Grassmannian) of the classical Darboux transformation of an n -th Gelfand–Dickey Lax operator L^n .

The classical Darboux transformation consists of factorizing an n -th order differential operator $Q = RS$, with R and S differential operators of order r , s , respectively, with $r + s = n$, and then exchanging the place of the factors, i.e., defining a new differential operator $Q' = SR$. We assume that $Q = L^n$ is a Gelfand–Dickey operator and we want

that Q' is again a Gelfand–Dickey operator. Let $L = P_W \partial P_W^{-1}$, with $W \in Gr(H)$, since $Q' = SQS^{-1}$, we find that $Q' = L'^n$, with $L' = P_{W'} \partial P_{W'}^{-1}$ and $P_{W'} = SP_W$. Because $Q = RS$ we find that

$$z^n W \subset W' \subset W,$$

hence the vectors $w_1(z), w_2(z), \dots, w_s(z) \in W$, that define the Bäcklund–Darboux transformation S , must be perpendicular to $z^n W$ and thus must belong to U , defined by (8.2). We thus have proven

Theorem 8.2 *Let $W \in Gr(H)$ satisfy $z^n W \subset W$ and let $L = P_W \partial P_W^{-1}$. The classical Darboux transformation of the Gelfand–Dickey differential operator $Q = L^n$, consists of choosing s ($0 < s < n$) linearly independent vectors $v_j \in U$ (U defined by (8.2)), such that*

$$S = \mathcal{W}(\psi_W; w_1, w_2, \dots, w_s)^{-1} \mathcal{W}(\psi_W; w_1, w_2, \dots, w_s; \partial).$$

Then $Q' = SQS^{-1}$ is again a Gelfand–Dickey differential operator.

9 From KP to the 1-Toda Lattice Hierarchy

The 1-Toda lattice hierarchy, as described in [1] (see also [3] and [11]) is the set of deformation equations (Lax equations)

$$\frac{\partial L}{\partial_n} = [(L^n)_+, L], \quad (9.1)$$

for infinite matrices

$$L = L(t, \Lambda) = \Lambda + \sum_{i=0}^{\infty} a_i \Lambda^{-i} \quad \text{with } \Lambda = \sum_{k \in \mathbb{Z}} \frac{\epsilon_k}{\epsilon_{k+1}} E_{k, k+1}, \quad (9.2)$$

depending on $t = (t_1, t_2, \dots)$ where $\partial_n = \partial / \partial t_n$ and where $\epsilon_k \in \mathbb{C}^\times$, $a_i = \sum_{k \in \mathbb{Z}} a_i(k, t) E_{kk}$ is an infinite diagonal matrix. Here A_+ refers to the upper-diagonal part of the matrix A , including the diagonal. We denote by $A_- = A - A_+$. For notational convenience we write $\Lambda = \epsilon \sum_{k \in \mathbb{Z}} E_{k, k+1} \epsilon^{-1}$ with $\epsilon = \sum_{k \in \mathbb{Z}} \epsilon_k E_{kk}$.

To the problem (9.1) one associates wavevectors $\Psi(t, z)$ and adjoint wavevectors $\Psi^*(t, z)$,

$$\begin{aligned} \Psi(t, z) &= (\epsilon_n \psi_n(t, z))_{n \in \mathbb{Z}} = P(t, \Lambda) e^{\sum_{k=0}^{\infty} t_k \Lambda^k} \epsilon \chi(z), \\ \Psi^*(t, z) &= (\epsilon_n^{-1} z \psi_{n+1}^*(t, z))_{n \in \mathbb{Z}} = (P(t, \Lambda)^T)^{-1} e^{-\sum_{k=0}^{\infty} t_k \Lambda^{-k}} \epsilon^{-1} \chi(z^{-1}), \end{aligned} \quad (9.3)$$

which satisfies

$$\begin{aligned} L \Psi(t, z) &= z \Psi(t, z), & L^T \Psi^*(t, z) &= z \Psi^*(t, z), \\ \partial_n(\Psi(t, z)) &= (L^n)_+ \Psi(t, z), & \partial_n(\Psi^*(t, z)) &= ((L^n)_+)^T \Psi^*(t, z). \end{aligned} \quad (9.4)$$

Here A^T stands for the transposed of the matrix A , $\chi(z)$ is the infinite column-vector $\chi(z) = (z^n)_{n \in \mathbb{Z}}$ and

$$P(t, \Lambda) = I + \sum_{k=1}^{\infty} p_k \Lambda^{-k}. \quad (9.5)$$

Notice that in this situation $L = P(t, \Lambda) \Lambda P(t, \Lambda)^{-1}$. We can formulate the 1-Toda lattice hierarchy in an other way, viz. as follows (see also [1]):

Proposition 9.1 *The following two statements are equivalent:*

- (a) *The Lax operator L of the form (9.2) satisfies the 1-Toda lattice hierarchy (9.1).*
 (b) *$\psi_n(t, z)$ is a wavefunction of the KP hierarchy for all $n \in \mathbb{Z}$, with $\psi_n^*(t, z)$ its adjoint wavefunction. These (adjoint) wave functions satisfy the $n - m$ -th modified KP hierarchy:*

$$\langle \psi_n(t, z) | \overline{z \psi_m^*(s, z)} \rangle = 0 \quad \text{for all } n \geq m. \quad (9.6)$$

To prove this proposition we use the following lemma:

Lemma 9.1 (a) *Let U and V be two infinite matrices, then*

$$\text{Res}_{z=0} \frac{1}{z} U \chi(z) (V \chi(z^{-1}))^T = UV^T.$$

(b) *Let $P(t, \partial)$ and $Q(t, \partial)$ be two pseudodifferential operators such that*

$$\text{Res}_{z=0} P(t, z) Q(t, -z) e^{\sum (t_i - s_i) z^i} = 0,$$

then $(P(t, \partial) Q(t, \partial)^)_- = 0$.*

Proof. For a proof of (b) we refer to [14]. To prove (a), let $U = \sum_{i,j} u_{ij} E_{ij}$ and $V = \sum_{i,j} v_{ij} E_{ij}$, then

$$\begin{aligned} \text{Res}_{z=0} \frac{1}{z} U \chi(z) (V \chi(z^{-1}))^T &= \text{Res}_{z=0} \frac{1}{z} \sum_{i,j,k,\ell} u_{ij} z^j v_{k\ell} z^{-\ell} E_{ik} \\ &= \sum_{i,j,k} u_{ij} v_{kj} E_{ik} \\ &= UV^T, \end{aligned}$$

which proves (a). □

Proof of Proposition 9.1. For all $k \geq 0$ one has

$$\begin{aligned} 0 &= (\Lambda^k)_- = (\Lambda^k P(t, \lambda) P(t, \lambda)^{-1})_- \\ &= (\Lambda^k P(t, \lambda) e^{\sum t^i \Lambda^i} e^{-\sum t^i \Lambda^i} P(t, \lambda)^{-1})_-, \end{aligned}$$

hence

$$\text{Res}_{z=0} \left(\frac{1}{z} \Lambda^k (\Psi(t, z)) \Psi^*(t, z)^T \right)_- = 0$$

and thus

$$\begin{aligned} 0 &= \text{Res}_{z=0} \left(\frac{1}{z} ((L^{n_1})_+)^{k_1} ((L^{n_2})_+)^{k_2} \dots ((L^{n_m})_+)^{k_m} (\Psi(t, z)) \Psi^*(t, z)^T \right)_- \\ &= \text{Res}_{z=0} \left(\frac{1}{z} \frac{\partial^{k_1 + \dots + k_m} (\Psi(t, z))}{\partial t_{n_1}^{k_1} \dots \partial t_{n_m}^{k_m}} \Psi^*(t, z)^T \right)_-. \end{aligned}$$

Now using Taylor's formula one obtains

$$\text{Res}_{z=0} \left(\frac{1}{z} \Psi(t, z) \Psi^*(s, z)^T \right)_- = 0, \quad (9.7)$$

which is equivalent to (9.6). Next we write the oscillating functions $\psi_n(t, z)$ and oscillating function $\psi^*(t, z)$ in terms of pseudodifferential operators.

$$\psi_n(t, z) = P(\partial)\partial^n e^{\sum t_i z^i} \quad \text{and} \quad \psi_n^*(t, z) = Q(\partial)(-\partial)^{-n} e^{-\sum t_i z^i}.$$

Using Lemma 9.1(b), one deduces from (9.6) for $n = m$ that $Q^* = P^{-1}$ and hence $\psi_n^*(t, z)$ is the adjoint oscillating function of $\psi_n(t, z)$. Since they satisfy (9.6) for $n = m$, $\psi_n(t, z)$ (resp. $\psi_n^*(t, z)$) must be a KP wavefunction (resp. adjoint wavefunction).

To prove the converse we use the bilinear identity (9.7) which is equivalent to (9.6). Denote

$$\Psi(t, z) = P(t, \Lambda) e^{\sum t_i \Lambda^i} \epsilon \chi(z) \quad \text{and} \quad \Psi^*(t, z) = Q(t, \Lambda) e^{-\sum t_i \Lambda^{-i}} \epsilon^{-1} \chi(z^{-1}),$$

then using part (a) of Lemma 9.1, we deduce from (9.7) with $s = t$ that $(P(t, \Lambda)Q(t, \Lambda)^T)_- = 0$ and hence

$$Q(t, \Lambda)^T = P(t, \Lambda)^{-1}. \quad (9.8)$$

Next differentiate (9.7) to t_n , then

$$\text{Res}_{z=0} \left(\frac{1}{z} (\partial_n(P(t, \Lambda)) + P(t, \Lambda)\Lambda^n) e^{\sum t_i \Lambda^i} \epsilon \chi(z) \Psi^*(s, z)^T \right)_- = 0.$$

Putting $s = t$ we deduce from this, using (9.8) and the Lemma 9.1 that

$$\partial_n(P(t, \Lambda))P(t, \Lambda)^{-1} = -(P(t, \Lambda)\Lambda^n P(t, \Lambda)^{-1})_-,$$

which is the Sato–Wilson equation for the 1-Toda lattice hierarchy. Differentiate $L(t, \Lambda) = P(t, \Lambda)\Lambda P(t, \Lambda)^{-1}$ to t_n , then using this Sato–Wilson equation one shows easily that L satisfies (9.1). $\square$

As a consequence of the proof of Proposition 9.1 one has the Sato–Wilson equation

$$\partial_n(P(t, \Lambda))P(t, \Lambda)^{-1} = -(P(t, \Lambda)\Lambda^n P(t, \Lambda)^{-1})_-. \quad (9.9)$$

Now return to the Segal–Wilson setting, then part (b) of the above proposition tells us that the planes $W_n \in Gr^{(n)}(H)$ related to ψ_n form an infinite flag, in fact one has two infinite flags

$$\begin{aligned} \cdots \subset W_{n+1} \subset W_n \subset W_{n-1} \subset W_{n-2} \subset \cdots, \\ \cdots \supset W_{n+1}^\perp \supset W_n^\perp \supset W_{n-1}^\perp \supset W_{n-2}^\perp \supset \cdots. \end{aligned} \quad (9.10)$$

So starting with one plane $W \in Gr^{(n)}(H)$, with wavefunction $\psi_W(t, z)$, adjoint wavefunction $\psi_W^*(t, z)$ and tau-function τ_W , we can apply recursively Bäcklund–Darboux transformations and adjoint Bäcklund–Darboux transformations to construct a whole flag of spaces. To be more precise:

Proposition 9.2 *Let $W \in Gr^{(n)}(H)$, with wavefunction $\psi_W(t, z)$, adjoint wavefunction $\psi_W^*(t, z)$ and tau-function τ_W and let $w_n, w_{n+1}, w_{n+2}, \dots$ be an ordered Hilbert basis of W and let $w_{n-1}, w_{n-2}, w_{n-3}, \dots$ be an ordered Hilbert basis of $W^\perp$, then the spaces*

$$W_m = \begin{cases} \{w \in W \mid \langle w | w_k \rangle = 0 \text{ for } k < m\} & \text{for } m > n, \\ W & \text{if } m = n, \\ W \oplus \sum_{k=m}^{n-1} \mathbb{C} w_k & \text{for } m < n \end{cases} \quad (9.11)$$

and spaces $W_m^\perp$ form infinite flags of the form (9.10). The corresponding wavefunctions $\psi_m(t, z)$, adjoint wavefunction $\psi_m^*(t, z)$ and tau-function τ_m are equal to (cf. (7.23))

$$\psi_m = \begin{cases} \mathcal{W}(\psi_W; w_n, \dots, w_{m-1}; \psi_W^*; \emptyset)^{-1} \mathcal{W}(\psi_W; w_n, \dots, w_{m-1}; \psi_W^*; \emptyset; \psi_W) & \text{for } m > n, \\ \psi_W & \text{if } n = m, \\ \mathcal{W}(\psi_W; \emptyset; w_m^\dagger, \dots, w_{n-1}^\dagger)^{-1} \mathcal{W}(\psi_W; \emptyset; \psi_W^*; w_m^\dagger, \dots, w_{n-1}^\dagger; \psi_W) & \text{for } m < n \end{cases} \quad (9.12)$$

and

$$\psi_m^* = \begin{cases} (-)^{n+m+1} \mathcal{W}(\psi_W^*; \emptyset; \psi_W; w_n, \dots, w_{m-1})^{-1} \mathcal{W}(\psi_W^*; \emptyset; \psi_W; w_n, \dots, w_{m-1}; \psi_W^*) & \text{for } m > n, \\ \psi_W^* & \text{if } n = m, \\ (-)^{n+m+1} \mathcal{W}(\psi_W^*; w_m^\dagger, \dots, w_{n-1}^\dagger; \psi_W; \emptyset)^{-1} \mathcal{W}(\psi_W^*; w_m^\dagger, \dots, w_{n-1}^\dagger; \psi_W; \emptyset; \psi_W^*) & \text{for } m < n \end{cases} \quad (9.13)$$

$$\tau_m = \begin{cases} \mathcal{W}(\psi_W; w_n, \dots, w_{m-1}; \psi_W^*; \emptyset) \tau_W & \text{for } m > n, \\ \tau_W & \text{if } n = m, \\ \mathcal{W}(\psi_W; \emptyset; \psi_W^*; w_m^\dagger, \dots, w_{n-1}^\dagger) \tau_W & \text{for } m < n, \end{cases} \quad (9.14)$$

Moreover the vector $\Psi(t, z) = (\epsilon_n \psi_n(t, z))_{n \in \mathbb{Z}}$ is a wavevector and $\Psi^*(t, z) = (\epsilon_n^{-1} z \psi_{n+1}^*(t, z))_{n \in \mathbb{Z}}$ is its adjoint wavevector of the 1-Toda lattice hierarchy.

The proof of this proposition is straightforward.

10 Some Applications

In all the applications that we consider in this section, we start with $W = W_0 = H_+$, hence $\tau_0 = 1$, $\psi_0(t, z) = e^{\sum_i t_i z^i}$ and $\psi_0^*(t, z) = e^{-\sum_i t_i z^i}$. Let w_j , $0 \leq j \leq n-1$, be the following elements of H_+ :

$$w_j = \sum_{i=0}^{\infty} \bar{\xi}_{ij} z^i, \quad \text{with } \sum_{i=0}^{\infty} |\xi_{ij}|^2 < \infty. \quad (10.1)$$

Then for $e^{\sum_i t_i z^i} = \sum_{\ell \in \mathbb{Z}} p_\ell(t) z^\ell$ one finds the following tau-functions:

$$\begin{aligned} \tau_n &= \mathcal{W}(\psi_0; w_0, \dots, w_{n-1}; \psi_0^*; \emptyset) \\ &= \det \left(\left\langle \frac{\partial^i e^{\sum_k t_k z^k}}{\partial x^i} \middle| \sum_{\ell=0}^{\infty} \bar{\xi}_{\ell j} z^\ell \right\rangle_{0 \leq i, j < n} \right) \\ &= \det \left(\sum_{\ell=0}^{\infty} p_{\ell-i} \xi_{\ell j} \right)_{0 \leq i, j < n}. \end{aligned} \quad (10.2)$$

In a similar way we also find for

$$w_j = \sum_{i=1}^{\infty} \bar{\xi}_{-i-j} z^{-i} \in H_-, \quad \text{with } \sum_{i=1}^{\infty} |\xi_{-i-j}|^2 < \infty, \quad (10.3)$$

that

$$\tau_{-n} = \det \left(\sum_{\ell=1}^{\infty} p_{\ell-i} \xi_{-\ell-j} \right)_{1 \leq i, j \leq n} \quad (10.4)$$

is a KP tau-function.

Notice that if we let n go to infinity, (10.2) is a special case of the famous result of Sato (see e.g. [16]).

By choosing special elements w_j , one obtains special KP tau-functions. For instance

$$w_j = \sum_{k=0}^{N-1} \bar{\eta}_{kj} \sum_{i=0}^{\infty} (\bar{\lambda}_k z)^i$$

leads to Hirota's N soliton solution (see also [16]):

$$\tau_n = \sum_{0 \leq i_0 < i_1, \dots, i_{n-1} \leq N-1} e^{\sum_{k=0}^{n-1} \sum_{j=1}^{\infty} t_j \lambda_{i_k}^j} \Delta(\lambda_{i_0}, \dots, \lambda_{i_{n-1}}) \det(\eta_{ipq})_{0 \leq p, q < n},$$

where $\Delta(\lambda_{i_0}, \dots, \lambda_{i_{n-1}})$ is the Vandermonde determinant.

Now return to the expressions (10.2) and (10.4). Here we have shown a well-known result of Ohta, Satsuma, Takahashi and Tokihiro [20], viz that if $f_0, f_1, \dots, f_{n-1}$ (resp. $f_{-1}, f_{-2}, \dots, f_{-n}$) are functions satisfying:

$$\begin{aligned} \frac{\partial f_j}{\partial t_k} &= \frac{\partial^k f_j}{\partial x^k} & \text{for } j \geq 0 \text{ and} \\ \frac{\partial f_j}{\partial t_k} &= (-)^{k-1} \frac{\partial^k f_j}{\partial x^k} & \text{for } j < 0, \end{aligned}$$

then the Wronskian determinants

$$\mathcal{W}(f_0, f_1, \dots, f_{n-1}), \quad (\text{resp. } \mathcal{W}(f_{-1}, f_{-2}, \dots, f_{-n})) \quad (10.5)$$

are tau-functions of the KP hierarchy.

In the article [19], Oevel and Strampp determine when the Wronskian solutions of the form (10.5) belong to the so-called m -vector k -constrained KP hierarchy. This is a reduction of the KP hierarchy, generalizing the Gelfand-Dickey hierarchies. Here one assumes that the KP Lax operator L satisfies

$$L^k = (L^k)_+ + \sum_{j=1}^m q_j(t) \partial^{-1} r_j(t),$$

with $q_j(t)$ eigenfunctions and $r_j(t)$ adjoint eigenfunctions of L . The authors gave in [12], [13] a geometrical interpretation of this m -vector k -constrained KP hierarchy in the Segal-Wilson setting. They showed that $L = L_W$ belongs to this m -vector k -constrained KP hierarchy if and only if the corresponding $W \in Gr(H)$ contains a subspace W' of codimension m such that $z^k W' \subset W$. We will now use this result to prove geometrically the following result of Oevel and Strampp [19]:

Proposition 10.1 *the tau functions $\mathcal{W}(f_0, f_1, \dots, f_{n-1})$ (resp. $\mathcal{W}(f_{-1}, f_{-2}, \dots, f_{-n})$) belong to the m -vector k -constrained KP hierarchy if and only if*

$$\begin{aligned} & \mathcal{W}(f_0, f_1, \dots, f_{n-1}, \frac{\partial^k f_{i_1}}{\partial x^k}, \dots, \frac{\partial^k f_{i_{m+1}}}{\partial x^k}) = 0, \\ & (\text{resp. } \mathcal{W}(f_{-1}, f_{-2}, \dots, f_{-n}) \frac{\partial^k f_{i_1}}{\partial x^k}, \dots, \frac{\partial^k f_{i_{m+1}}}{\partial x^k}) = 0), \end{aligned}$$

for all $(m+1)$ -tuples $(f_{i_1}, \dots, f_{i_{m+1}})$ with indices $0 \leq i_1 < i_2 < \dots < i_{m+1} \leq n-1$ (resp. $-n \leq i_1 < i_2 < \dots < i_{m+1} \leq -1$).

Proof. First consider the positive case, viz $n \geq 0$. Write $f_j = \langle e^{\sum_k t_k z^k} | w_j \rangle$ with w_j as in (10.1). Then

$$\tau_n = \mathcal{W}(f_0, f_1, \dots, f_{n-1}) = \mathcal{W}(\psi_0; w_0, \dots, w_{n-1}; \psi_0^*; \emptyset).$$

If this is a solution of the m -vector k -constrained KP hierarchy, then the space

$$W_n = \{v \in H_+ | \langle v | w_i \rangle = 0 \text{ for } 0 \leq i \leq n-1\}$$

has a subspace W' of codimension m such that $z^k W' \subset W$. In other words, one has that for all $w \in W'$ and $0 \leq j \leq n-1$ both $\langle w | w_j \rangle = 0$ and $\langle z^k w | w_j \rangle = \langle w | z^{-k} w_j \rangle = 0$. Which means that

$$W' \subset \{v \in H_+ | \langle v | w_i \rangle = \langle v | z^{-k} w_i \rangle = 0 \text{ for } 0 \leq i \leq n-1\} \subset W.$$

or equivalently that

$$\dim \langle w_0, w_1, \dots, w_{n-1}, (z^{-k} w_0)_+, (z^{-k} w_1)_+, \dots, (z^{-k} w_{n-1})_+ \rangle \leq n + m,$$

where $u_+ = \sum_{k=0}^{\infty} u_k z^k$ if $u = \sum_{k \in \mathbb{Z}} u_k z^k$. In other words

$$\begin{aligned} & \mathcal{W}(f_0, f_1, \dots, f_{n-1}, \frac{\partial^k f_{i_1}}{\partial x^k}, \dots, \frac{\partial^k f_{i_{m+1}}}{\partial x^k}) = \\ & \mathcal{W}(\psi_0; w_0, \dots, w_{n-1}, z^{-k} w_{i_1}, \dots, z^{-k} w_{i_{m+1}}; \psi_0^*; \emptyset) = 0, \end{aligned}$$

for all $0 \leq i_1 < i_2 < \dots < i_{m+1} \leq n-1$.

In the negative case, one writes $f_j = \langle e^{-\sum_k t_k z^k} | w_j \rangle$ with w_j as in (10.3). One then has to notice that the m -vector k -constrained KP hierarchy means that W is a subspace of codimension m of W'' such that $z^k W \subset W''$ or equivalently $z^{-k} (W'')^\perp \subset W^\perp$ and one can finish the proof along the same lines as in the positive case. $\square$

The next application generates certain orthogonal polynomials. It is related to Matrix models, see e.g [3], [11] and references therein.

Let (a, b) be an interval of $\mathbb{R}$ and $w(y)$ a weight function which is non-negative. With these data we associate the inner product

$$(f_1 | f_2) = \int_a^b f_1(y) f_2(y) w(y) dy, \quad (10.6)$$

which is well defined for all functions f for which $\sqrt{w}f$ is quadratically integrable on (a, b) .

The numbers

$$c_n = \int_a^b y^n w(y) dy, \quad (10.7)$$

are called the *moments* of the weight function w . It is well-known (see e.g. [6]) that the polynomials

$$p_n(y) = \frac{\det \begin{pmatrix} c_0 & c_1 & \cdots & c_n \\ c_1 & c_2 & \cdots & c_{n+1} \\ \vdots & \vdots & \dots & \vdots \\ c_{n-1} & c_n & \cdots & c_{2n-1} \\ 1 & y & \cdots & y^n \end{pmatrix}}{\det(c_{i+j})_{0 \leq i, j \leq n-1}} \quad (10.8)$$

form an orthogonal set, i.e.,

$$(p_m | p_n) = \delta_{mn} \frac{\det(c_{i+j})_{0 \leq i, j \leq n}}{\det(c_{i+j})_{0 \leq i, j \leq n-1}}.$$

Now consider the generating series of the complex conjugates of the moments

$$f(z) = \sum_{k=0}^{\infty} \bar{c}_k z^k. \quad (10.9)$$

Since the moments of the weight function w satisfy the inequality

$$|c_n| \leq \max(|a|, |b|)^n M,$$

with M a constant, the series $f(z)$ converges on a neighborhood of zero. Hence there is a $\lambda_0 > 0$ such that $\sum_{n \geq 0} |c_n|^2 \lambda_0^{2n} < \infty$. For all λ with $0 < \lambda \leq \lambda_0$ we define the element $f_i^{(\lambda)}(z) \in H$ by

$$f_i^{(\lambda)}(z) = \sum_{n=0}^{\infty} \bar{c}_n \lambda^n z^{n-i}, \quad \text{for } i = 0, 1, 2, \dots \quad (10.10)$$

Now we consider the following chain of subspaces of H_+ :

$$W_0^{(\lambda)} = H_+ \quad \text{and} \quad W_n^{(\lambda)} = \{w \in H_+ \mid \langle w | f_i^{(\lambda)} \rangle = 0 \text{ for } i = 0, 1, \dots, n-1\}.$$

Note that one can find for each $k \geq 1$ a polynomial $q_k^{(\lambda)}$ of degree k such that $\langle q_k^{(\lambda)} | f \rangle = 0$. Since the topological span of the $\{q_k^{(\lambda)} \mid k \geq 1\}$ has codimension one in H_+ and the same holds for $W_1^{(\lambda)}$, we must have

$$W_1^{(\lambda)} = \text{the closure of } \text{Span}\{q_k^{(\lambda)} \mid k \geq 1\}.$$

Now we like to choose the $\{q_k^{(\lambda)}\}$ such that for all $m \geq 1$

$$W_m^{(\lambda)} = \text{the closure of } \text{Span}\{q_k^{(\lambda)} \mid k \geq m\}.$$

Then the whole chain of subspaces $\{W_m^{(\lambda)} | m \geq 1\}$ has been expressed in the sequence of polynomials $\{q_k^{(\lambda)} | k \geq 1\}$. Because of the codimension one argument, it is necessary and sufficient that for all $m \geq 1$ and all $j \geq 0$, $j < m$,

$$\langle q_m^{(\lambda)} | f_j^{(\lambda)} \rangle = 0.$$

This condition will tell you how to make an appropriate choice for the $\{q_k^{(\lambda)} | k \geq 1\}$. For if $q_m^{(\lambda)}(y) = \sum_{\ell=0}^m a_{m\ell} y^\ell$, then we have

$$\begin{aligned} \langle q_m^{(\lambda)} | f_j^{(\lambda)} \rangle &= \langle \sum_{\ell=0}^m a_{m\ell} y^\ell | \sum_{r=0}^m \bar{c}_r \lambda^r z^{r-j} \rangle \\ &= \sum_{\ell=0}^m a_{m\ell} c_{\ell+j} \lambda^j \\ &= \sum_{\ell=0}^m \int_a^b a_{m\ell} y^{\ell+j} \lambda^j w(y) dy \\ &= \int_a^b q_m^{(\lambda)}(\lambda y) y^j w(y) dy \end{aligned} \tag{10.11}$$

Hence, if we choose $q_m^{(\lambda)}(\lambda y) = p_m(y)$, then due to $(y^j, p_m(y)) = 0$ for $j < m$, we get that (10.11) is equal to zero. Thus (cf. [11])

$$W_m^{(\lambda)} = \text{the closure of Span}\{p_k\left(\frac{z}{\lambda}\right) | k \geq m\}. \tag{10.12}$$

Next we compute the tau-function corresponding to this chain following (10.2). We consider $N > \max(|a|, |b|, 1)$ and the KP-flows from $\Gamma_+(N)$. In that case we have

$$\begin{aligned} \langle \frac{\partial^p e^{\sum_k t_k z^k}}{\partial x^p} | f_q^{(\lambda)}(z) \rangle &= \langle z^{p+q} e^{\sum_k t_k z^k} | f_0^{(\lambda)}(z) \rangle \\ &= \int_a^b (\lambda y)^{p+q} e^{\sum_k t_k (\lambda y)^k} w(y) dy \\ &= \lambda^{p+q} c_{p+q}^{(\lambda)}(t), \end{aligned}$$

where $c_n^{(\lambda)}(t)$ is the n -th moment of the t, λ -dependent inner product (cf. (10.7))

$$(f_1(t, z) | f_2(t, z))_t^{(\lambda)} = \int_a^b f_1(t, z) f_2(t, z) e^{\sum_k t_k (\lambda z)^k} w(z) dz. \tag{10.13}$$

Note that due to the fact that we have taken our flows from $\Gamma_+(N)$, this is a well-defined inner product for real t_i 's, even if $\lambda = 1$.

We find that the to $W_n^{(\lambda)}$ corresponding wave function $\psi_n^{(\lambda)}(t, z) = Q_n^{(\lambda)}(t, z) e^{\sum_k t_k z^k}$ is

equal to $BD(W_n^{(\lambda)}, H_+)e^{\sum_k t_k z^k}$. Hence,

$$\begin{aligned}
Q_n^{(\lambda)}(t, z) &= \frac{\det \begin{pmatrix} c_0^{(\lambda)}(t) & c_1^{(\lambda)}(t)\lambda & \cdots & c_n^{(\lambda)}(t)\lambda^n \\ c_1^{(\lambda)}(t)\lambda & c_2^{(\lambda)}(t)\lambda^2 & \cdots & c_{n+1}^{(\lambda)}(t)\lambda^{n+1} \\ \vdots & \vdots & \cdots & \vdots \\ c_{n-1}^{(\lambda)}(t)\lambda^{n-1} & c_n^{(\lambda)}(t)\lambda^n & \cdots & c_{2n-1}^{(\lambda)}(t)\lambda^{2n-1} \\ 1 & z & \cdots & z^n \end{pmatrix}}{\det(c_{i+j}^{(\lambda)}(t)\lambda^{i+j})_{0 \leq i, j \leq n-1}} \\
&= \frac{\det \left(\begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & \lambda^{n-1} & \\ & & & 1 \end{pmatrix} \begin{pmatrix} c_0^{(\lambda)}(t) & c_1^{(\lambda)}(t) & \cdots & c_n^{(\lambda)}(t) \\ c_1^{(\lambda)}(t) & c_2^{(\lambda)}(t) & \cdots & c_{n+1}^{(\lambda)}(t) \\ \vdots & \vdots & \cdots & \vdots \\ c_{n-1}^{(\lambda)}(t) & c_n^{(\lambda)}(t) & \cdots & c_{2n-1}^{(\lambda)}(t) \\ 1 & \frac{z}{\lambda} & \cdots & \left(\frac{z}{\lambda}\right)^n \end{pmatrix} \begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & & \lambda^n \end{pmatrix} \right)}{\lambda^{n(n-1)} \det(c_{i+j}^{(\lambda)}(t))_{0 \leq i, j \leq n-1}} \\
&= \lambda^n P_n^{(\lambda)}\left(t, \frac{z}{\lambda}\right),
\end{aligned} \tag{10.14}$$

where

$$P_n^{(\lambda)}(t, z) = \frac{\det \begin{pmatrix} c_0^{(\lambda)}(t) & c_1^{(\lambda)}(t) & \cdots & c_n^{(\lambda)}(t) \\ c_1^{(\lambda)}(t) & c_2^{(\lambda)}(t) & \cdots & c_{n+1}^{(\lambda)}(t) \\ \vdots & \vdots & \cdots & \vdots \\ c_{n-1}^{(\lambda)}(t) & c_n^{(\lambda)}(t) & \cdots & c_{2n-1}^{(\lambda)}(t) \\ 1 & z & \cdots & z^n \end{pmatrix}}{\det(c_{i+j}^{(\lambda)}(t))_{0 \leq i, j \leq n-1}}. \tag{10.15}$$

These $P_n^{(\lambda)}(t, z)$ form the orthogonal polynomials with respect to the t, λ -dependent inner product $(\cdot | \cdot)_t^{(\lambda)}$, defined by (10.13). Moreover, the $\psi_n^{(\lambda)}$'s for $n = 0, 1, 2, \dots$ form a set of 1-Toda wavefunctions and the $\tau_n^{(\lambda)}$'s or $\sigma_n^{(\lambda)}$'s where

$$\begin{aligned}
\tau_n^{(\lambda)}(t) &= \det(c_{i+j}^{(\lambda)}(t)\lambda^{i+j})_{0 \leq i, j \leq n-1} = \lambda^{n(n-1)} \sigma_n^{(\lambda)}(t), \\
\sigma_n^{(\lambda)}(t) &= \det(c_{i+j}^{(\lambda)}(t))_{0 \leq i, j \leq n-1},
\end{aligned} \tag{10.16}$$

form a set of 1-Toda tau-functions.

For all $w_n \in W_n^{(\lambda)}$ we have that $\langle w_n | f_i^{(\lambda)} \rangle = 0$ for $i = 0, 1, \dots, n-1$, then $\langle zw_n | f_i^{(\lambda)} \rangle = \langle w_n | z^{-1} f_i^{(\lambda)} \rangle = \langle w_n | f_{i+1}^{(\lambda)} \rangle$ is zero for $i = 0, 1, \dots, n-2$. Hence $zW_n^{(\lambda)} \subset W_{n-1}^{(\lambda)}$ and we deduce that $\psi_n^{(\lambda)}(t, z)$ belongs to the 1-vector 1-constrained KP hierarchy, i.e.,

$$L_{W_n^{(\lambda)}} = \partial + q_n^{(\lambda)} \partial^{-1} r_n^{(\lambda)},$$

with $q_n^{(\lambda)}$ and $r_n^{(\lambda)}$ eigenfunctions, respectively adjoint eigenfunctions of L_{W_n} . From the construction in Section 3 of [12], it is easy to determine the $q_n^{(\lambda)}$ and $r_n^{(\lambda)}$'s, in fact

$$L_{W_n^{(\lambda)}} = \partial + g_n \frac{\tau_{n+1}^{(\lambda)}}{\tau_n^{(\lambda)}} \partial^{-1} \frac{\tau_{n-1}^{(\lambda)}}{\tau_n^{(\lambda)}},$$

with $g_n \in \mathbb{C}^\times$. Hence

$$\begin{aligned} z\psi_n^{(\lambda)} &= \frac{\partial \psi_n^{(\lambda)}}{\partial x} + g_n \frac{\tau_{n+1}^{(\lambda)}}{\tau_n^{(\lambda)}} \partial^{-1} \frac{\tau_{n-1}^{(\lambda)}}{\tau_n^{(\lambda)}} (\psi_n^{(\lambda)}) \\ &= \frac{\partial \psi_n^{(\lambda)}}{\partial x} + g_n \frac{\tau_{n+1}^{(\lambda)}}{\tau_n^{(\lambda)}} \partial^{-1} \frac{\tau_{n-1}^{(\lambda)}}{\tau_n^{(\lambda)}} \frac{\tau_n^{(\lambda)}}{\tau_{n-1}^{(\lambda)}} \partial \frac{\tau_{n-1}^{(\lambda)}}{\tau_n^{(\lambda)}} (\psi_{n-1}^{(\lambda)}) \\ &= \frac{\partial \psi_n^{(\lambda)}}{\partial x} + g_n \frac{\tau_{n+1}^{(\lambda)} \tau_{n-1}^{(\lambda)}}{(\tau_n^{(\lambda)})^2} \psi_{n-1}^{(\lambda)}. \end{aligned}$$

Since also

$$\psi_{n+1}^{(\lambda)} = \frac{\partial \psi_n^{(\lambda)}}{\partial x} - \frac{\partial}{\partial x} \left(\log \frac{\tau_{n+1}^{(\lambda)}}{\tau_n^{(\lambda)}} \right) \psi_n^{(\lambda)},$$

we find that

$$z\psi_n^{(\lambda)} = \psi_{n+1}^{(\lambda)} + \frac{\partial}{\partial x} \left(\log \frac{\tau_{n+1}^{(\lambda)}}{\tau_n^{(\lambda)}} \right) \psi_n^{(\lambda)} + g_n \frac{\tau_{n+1}^{(\lambda)} \tau_{n-1}^{(\lambda)}}{(\tau_n^{(\lambda)})^2} \psi_{n-1}^{(\lambda)}. \quad (10.17)$$

We can now determine the constant g_n . For this purpose we take the innerproduct $\langle z\psi_n^{(\lambda)} | f_{n-1}^{(\lambda)} \rangle$. Then from (10.17) we deduce that

$$g_n = \frac{\langle \psi_n^{(\lambda)} | f_n^{(\lambda)} \rangle \tau_n^{(\lambda)} \tau_n^{(\lambda)}}{\langle \psi_{n-1}^{(\lambda)} | f_{n-1}^{(\lambda)} \rangle \tau_{n-1}^{(\lambda)} \tau_{n+1}^{(\lambda)}} = \frac{\tau_{n+1}^{(\lambda)} \tau_n^{(\lambda)}}{\tau_n^{(\lambda)} \tau_{n+1}^{(\lambda)}} = 1$$

Thus we find that

$$\frac{z}{\lambda} P_n^{(\lambda)}(t, \frac{z}{\lambda}) = P_{n+1}^{(\lambda)}(t, \frac{z}{\lambda}) + \frac{1}{\lambda} \frac{\partial}{\partial x} \left(\log \frac{\sigma_{n+1}^{(\lambda)}}{\sigma_n^{(\lambda)}} \right) P_n^{(\lambda)}(t, \frac{z}{\lambda}) + \frac{\sigma_{n+1}^{(\lambda)} \sigma_{n-1}^{(\lambda)}}{(\sigma_n^{(\lambda)})^2} P_{n-1}^{(\lambda)}(t, \frac{z}{\lambda}). \quad (10.18)$$

This defines the recursion relation for the orthogonal polynomials $P_n^{(\lambda)}(t, z)$. Notice that the innerproduct (10.13), the formula's (10.14), (10.15), (10.16) and (10.18) are also well-defined for $\lambda = 1$. Hence let $P_n(t, z) = P_n^{(1)}(t, z) = Q_n^{(1)}(t, z)$ and $\tau_n = \tau_n^{(1)} = \sigma_n^{(1)}$, then the orthogonal polynomials $P_n(t, z)$, orthogonal with respect to the inner product $(\cdot | \cdot)_t^{(1)}$, satisfy the following recursion relation:

$$zP_n(t, z) = P_{n+1}(t, z) + \frac{\partial}{\partial x} \left(\log \frac{\tau_{n+1}}{\tau_n} \right) P_n(t, z) + \frac{\tau_{n+1} \tau_{n-1}}{\tau_n^2} P_{n-1}(t, z). \quad (10.19)$$

11 Appendix: Alternative Proof of Theorem 5.1

We want to give an alternative proof for expression of τ_{W_q} in Theorem 5.1. Let τ_W be the tau-function corresponding to $W \in Gr^{(\ell)}(H)$, then τ_W satisfies the bilinear identity

$$Res_{z=0} \tau_W(t - [z^{-1}]) e^{\sum_{i=1}^{\infty} (t_i - s_i) z^i} \tau_W(s + [z^{-1}]) = 0. \quad (11.1)$$

Now let the following operator

$$w^{2\ell} e^{\sum_{i=1}^{\infty} (t_i + s_i) w^i} e^{-\sum_{i=1}^{\infty} (\frac{\partial}{\partial t_i} + \frac{\partial}{\partial s_i}) \frac{w^{-i}}{i}}$$

act on this bilinear identity (11.1). Then we obtain after some calculation that

$$Res_{z=0} w^{2\ell} \tau_W(t - [z^{-1}] - [w^{-1}]) e^{\sum (t_i - \frac{z^i}{i}) w^i} e^{\sum (t_i - s_i) z^i} w^\ell \tau_W(s - [w^{-1}] + [z^{-1}]) e^{\sum (s_i + \frac{z^i}{i}) w^i} = 0$$

and hence

$$\tilde{\tau} := \psi_W(t, w) \tau_w(t) = w^\ell \tau_W(t - [w^{-1}]) e^{\sum_{i=1}^\infty t_i w^i}$$

is again a tau-function, however not a Segal–Wilson tau-function. Let $\tilde{\psi}(t, z)$ be the to $\tilde{\tau}$ corresponding wavefunction, i.e.,

$$\tilde{\psi}(t, z) = z^{\ell+1} \frac{\tau_W(t - [z^{-1}] - [w^{-1}]) e^{\sum_{i=1}^\infty (t_i - \frac{z^i}{i}) w^i}}{\tau_W(t - [w^{-1}]) e^{\sum_{i=1}^\infty t_i w^i}} e^{\sum_{i=1}^\infty t_i z^i},$$

then

$$\psi(t, w) \tilde{\psi}(t, z) = z \psi_W(t - [z^{-1}], w) \psi_W(t, z).$$

On the other hand $\tilde{\psi}(t, z)$ can be obtained with the Bäcklund–Darboux transformation

$$\partial - \frac{\partial \log(\tilde{\tau}/\tau)}{\partial x} = \partial - \psi_W(t, w)^{-1} \frac{\partial \psi_W(t, w)}{\partial x}$$

from $\psi_W(t, z)$. So we obtain that

$$\begin{aligned} \text{Wronskian}(\psi_W(t, z), \psi_W(t, w)) &= \frac{\partial \psi_W(t, z)}{\partial x} \psi_W(t, w) - \frac{\partial \psi_W(t, w)}{\partial x} \psi_W(t, z) \\ &= z \psi_W(t - [z^{-1}], w) \psi_W(t, z). \end{aligned}$$

Next let $\psi_q(t, z) = q(t) \partial q(t)^{-1} \psi_W(t, z)$ with $q(t) = \langle \psi_W(t, w) | s(w) \rangle$, then

$$\begin{aligned} q(t) \psi_q(t, z) &= \langle \psi_W(t, w) | s(w) \rangle \psi_q(t, z) \\ &= \langle \psi_W(t, w) | s(w) \rangle \frac{\partial \psi_W(t, z)}{\partial x} - \langle \frac{\partial \psi_W(t, w)}{\partial x} | s(w) \rangle \psi_W(t, z) \\ &= \langle \text{Wronskian}(\psi_W(t, z), \psi_W(t, w)) | s(w) \rangle \\ &= \langle z \psi_W(t - [z^{-1}], w) \psi_W(t, z) | s(w) \rangle \\ &= z q(t - [z^{-1}]) \psi_W(t, z), \end{aligned}$$

and

$$\psi_q(t, z) = z \frac{q(t - [z^{-1}])}{q(t)} \psi_W(t, z). \quad (11.2)$$

Now use formula (1.14) and one obtains that

$$\partial_k \log \frac{\tau_{W_q}(t)}{q(t) \tau_W(t)} = 0 \quad \text{for all } k = 1, 2, \dots$$

Thus we conclude that upto multiplication by a constant

$$\tau_{W_q}(t) = q(t) \tau_W(t). \quad (11.3)$$

The formula $\tau_{W_r}(t) = r(t) \tau_W(t)$ can be obtained in a similar matter.

References

- [1] M. Adler, E. Horozov, P. van Moerbeke, The solution to the q -KdV equation, preprint (1997)
- [2] M. Adler, P. van Moerbeke, Birkhoff Strata, Bäcklund Transformations and Regularization of Isospectral Operators, *Adv. in Math.* 108 (1994), 140-204)
- [3] M. Adler, P. van Moerbeke, Matrix integrals, Toda symmetries, Virasoro constraints and orthogonal polynomials, *Duke Math. J.* 80 (1995) 23-88.
- [4] H. Aratyn, E. Nissimov and S. Pacheva, Method of squared eigenfunction potentials in integrable hierarchies of KP type, *Commun. Math. Phys.* 193 (1998), 493– (solv-int 9701017).
- [5] B. Bakalov, E. Horozov, M. Yakimov, Bäcklund–Darboux transformations in Sat’s Grassmannian, *Serdica Math. J.* **22** (1996), 571–586.
- [6] H. Bateman, Higher transcendental functions, Vol II, McCraw-Hill, New York, 1953.
- [7] L.-L. Chau, J.C. Shaw, H.C. Yen, Solving the KP hierarchy by gauge transformations, *Commun. Math. Phys.* 149 (1992), 263-278.
- [8] M.M. Crum, Associated Sturm-Liouville systems, *Quart. J. Math.* 6 (1955), 121-127.
- [9] E. Date, M. Jimbo, M.Kashiwara, T. Miwa, Transformation groups for soliton equations, in: *Nonlinear integral systems – classical theory and quantum theory* eds. M. Jimbo, and T. Miwa, World Scientific, 39–120, 1983.
- [10] G.F. Helminck, G.F. Post, Geometrical interpretation of the bilinear equations for the KP hierarchy, *Letters Math. Phys.* **16** (1988), 359–364.
- [11] L. Haine, E. Horozov, Toda orbits of Laguerre polynomials and representations of the Virasoro algebra, *Bull. Sci. Math.* **117** (1993), 485–518.
- [12] G.F. Helminck, J.W. van de Leur, An analytic description of the vector constrained KP hierarchy, *Commun. Math Phys.* 193 (1998), 627–641 (solv-int 9706004)
- [13] G.F. Helminck, J.W. van de Leur. Constrained and Rational Reductions of the KP hierarchy. Preprint, to appear in "Supersymmetry and Integrable Models", Springer Lecture Notes in Physics 502, 1998, 167–182.
- [14] V.G. Kac, J.W. van de Leur. The n -component KP hierarchy and representation theory, in *Important developments in soliton theory*, eds. A.S. Fokas and V. E. Zakharov, Springer Series in Nonlinear Dynamics (1993), 302 – 343. Extended version of this paper with the same name will appear in the second edition of this book.
- [15] V.B. Matveev, M.A. Salle, Darboux Transformations and Solitons, Springer Series in Nonlinear Dynamics, Springer, Berlin, 1991.
- [16] M. Mulase, Algebraic theory of the KP equations, in *Interface between Mathematics and physics*, Ed. by S.T. Yau, International Press Company (1994), 157-223.

- [17] W. Oevel, Darboux theorems and Wronskian formulas for integrable systems (I. Constrained KP flows), *Physica A* 195 (1993), 533-576.
- [18] W. Oevel, W. Schief, Darboux theorems and the KP hierarchy, in: *Applications of analytic and geometric Methods in Differential Equations* (Proceedings of the NATO Advanced Research Workshop, Exeter, 14-17 July 1992, UK) P.A. Clarkson (ed.), Kluwer Publ. (1993), 193-206.
- [19] W. Oevel, W. Strampp, Wronskian solutions of the constrained Kadomtsev-Petviashvili hierarchy, *Journ. Math. Phys.* 37 (1996) p. 6213-6219.
- [20] Y. Ohta, J. Satsuma, D. Takahashi, T. Tokihiro, *Prog. Teor. Phys. Suppl.* 94 (1988), 219-
- [21] A. Pressley, G. Segal, *Loop groups*, Oxford Mathematical Monographs, Oxford University Press, Oxford 1988.
- [22] G. Segal and G. Wilson, Loop groups and equations of KdV type, *Publ. Math IHES* 63 (1985), p. 1-64.